

From THE DEPARTMENT OF  
MEDICAL BIOCHEMISTRY AND BIOPHYSICS  
Karolinska Institutet, Stockholm, Sweden

# **VEGF-B IN ISLETS OF LANGERHANS: ROLE IN VASCULAR SIGNALING AND LIPID HANDLING**

Frank Chenfei Ning, M.D.



**Karolinska  
Institutet**

Stockholm 2020

**Cover image:** An islet of Langerhans stained with insulin (green), glucagon (red) and vessel marker CD31 (white).

Image produced by Frank Chenfei Ning, 2018

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by E-Print AB

© Frank Chenfei Ning, 2020

ISBN 978-91-7831-743-1

# VEGF-B IN ISLETS OF LANGERHANS: ROLE IN VASCULAR SIGNALING AND LIPID HANDLING

## THESIS FOR DOCTORAL DEGREE (Ph.D.)

Public Defense at Karolinska Institutet

**Lecture Hall Inghesalen**, Widerströmska Huset, Tomtebodavägen 18A, Solna

**Friday 6th March, 2020 at 9:30**

By

**Frank Chenfei Ning**

*Principal Supervisor:*

Dr. Daniel Nyqvist  
Karolinska Institutet  
Department of Medical Biochemistry and Biophysics  
Division of Vascular Biology

*Co-supervisor(s):*

Dr. Ingrid Nilsson  
Karolinska Institutet  
Department of Medical Biochemistry and Biophysics  
Division of Vascular Biology

Professor Ulf Eriksson  
Karolinska Institutet  
Department of Medical Biochemistry and Biophysics  
Division of Vascular Biology

*Opponent:*

Professor Peter Bergsten  
Uppsala University  
Department of Medical Cell Biology

*Examination Board:*

Docent Peetra Magnusson  
Uppsala University  
Department of Immunology, Genetics and Pathology

Docent Mats Hellström  
Uppsala University  
Department of Immunology, Genetics and Pathology

Docent Christina Bark  
Karolinska Institutet  
Department of Molecular Medicine and Surgery





谨以此书献给敬爱的奶奶  
并谨此纪念我敬爱的爷爷

*To my dear grandmother and  
In loving memory of my grandfather*



## ABSTRACT

The discovery of Vascular endothelial growth factor (VEGF)-B's role in endothelial fatty acid uptake and tissue lipid accumulation opened a new arena for exploration of VEGF-B in lipid metabolism. Indeed, later studies demonstrated the beneficial/therapeutic effects of reducing VEGF-B signaling systemically. However, the tissue specific role of VEGF-B signaling has yet to be determined and furthermore the contribution of tissue specific VEGF-B activity to the systemic effects is also unknown to date.

In **paper I**, we investigated the role of local VEGF-B signaling in pancreatic islets by employing a novel mouse model where *Vegfb* in pancreatic  $\beta$ -cells were selectively deleted using Cre/loxP technology. We showed that *Vegfb* is abundantly expressed in both endocrine and exocrine pancreas, but that ablation of *Vegfb* from  $\beta$ -cells does not affect systemic glucose homeostasis or islet lipid uptake under chow or high-fat diet (HFD) conditions. However, deletion of *Vegfb* in pancreatic  $\beta$ -cells increased insulin (*Ins2*) gene expression, which indicates the potential interaction between VEGF-B signaling and transcriptional regulation of insulin, a phenotype warranting further studies.

In **paper II**, we continued to explore the role of pancreatic  $\beta$ -cell derived VEGF-B in systemic homeostasis. In contrast to the mouse model used in paper I, we utilized a mouse model where the human VEGF-B<sub>167</sub> isoform is overexpressed in pancreatic  $\beta$ -cells (under the rat insulin promoter). In this study, we could demonstrate that overexpression of VEGF-B in pancreatic  $\beta$ -cells does not affect systemic glucose homeostasis under chow or HFD conditions. Unexpectedly, overexpression of VEGF-B in pancreatic islets led to increased plasma triglycerides. This could indicate VEGF-B's involvement in plasma lipoprotein synthesis or degradation, and further studies are needed to elucidate this effect.

The role of VEGF-B in angiogenesis is somewhat ambiguous. To address this question, in **paper III**, we deployed a newly developed intravital imaging protocol where isolated mouse islets were transplanted into the anterior chamber of the eye to facilitate the observation of islet vessel development, pharmacological treatment of tumor islets and effects of VEGF-B overexpression in  $\beta$ -cells. We showed that the advantage of this platform comprises high spatial resolution, real-time monitoring of islet development and feasibility for evaluating efficacy of pharmacological treatments of tumor islets. In addition, we observed that

overexpression of VEGF-B in tumor islets increased tumor angiogenesis but impaired tumor growth, providing additional information about the role of VEGF-B in pathological angiogenesis.

Canonical Wnt/ $\beta$ -catenin signaling is essential for the retinal and cerebral angiogenesis during development. However, little is known regarding the contribution of endothelial  $\beta$ -catenin signaling in postnatal central nervous system (CNS) and potential interaction with other signaling pathways regulating angiogenesis. In **paper IV**, we studied the role of endothelial  $\beta$ -catenin signaling in central nervous system angiogenesis. We demonstrated that impaired Wnt/ $\beta$ -catenin signaling resulted in reduced postnatal retinal- and cerebral angiogenesis, presumably by diminished VEGFR2 expression. We furthermore identified crosstalk between the Wnt/ $\beta$ -catenin and Notch/VEGFA signaling pathways.

In summary, this thesis provides further knowledge about the specific role of VEGF-B signaling in islets of Langerhans, and its effects on systemic glucose homeostasis and lipid handling. Additionally, development of a novel intravital imaging protocol, as well as uncovering of additional roles of endothelial Wnt/ $\beta$ -catenin signaling in CNS, constitute parts of this thesis.

## LIST OF SCIENTIFIC PAPERS

- I. **Frank Chenfei Ning**, Nina Jensen, Jiarui Mi, William Lindström, Mirela Balan, Lars Muhl, Ulf Eriksson, Ingrid Nilsson & Daniel Nyqvist.  
VEGF-B ablation in pancreatic  $\beta$ -cells upregulates insulin expression without affecting glucose homeostasis or islet lipid uptake.  
*Scientific Reports* 10, 923 (2020) doi.org/10.1038/s41598-020-57599-2
- II. **Frank Chenfei Ning**, Mirela Balan, Jiarui Mi, Ulf Eriksson, Ingrid Nilsson & Daniel Nyqvist.  
Overexpression of VEGF-B in pancreatic  $\beta$ -cells does not affect glucose homeostasis but increases plasma triglycerides.  
*Manuscript*
- III. Mirela Balan, Marta Trusohamn, **Frank Chenfei Ning**, Stefan Jacob, Kristian Pietras, Ulf Eriksson, Per-Olof Berggren & Daniel Nyqvist.  
Noninvasive intravital high-resolution imaging of pancreatic neuroendocrine tumours.  
*Scientific Reports* 9, 14636 (2019) doi:10.1038/s41598-019-51093-0
- IV. Agnieszka Martowicz, Marta Trusohamn, Nina Jensen, Joanna Wisniewska-Kruk, Monica Corada, **Frank Chenfei Ning**, Julianna Kele, Elisabetta Dejana & Daniel Nyqvist.  
Endothelial  $\beta$ -Catenin Signaling Supports Postnatal Brain and Retinal Angiogenesis by Promoting Sprouting, Tip Cell Formation, and VEGFR (Vascular Endothelial Growth Factor Receptor) 2 Expression.  
*Arteriosclerosis, Thrombosis, and Vascular Biology*, 39, 2273-2288 (2019).  
doi:10.1161/ATVBAHA.119.312749

医，以人为本，以德为尚，以术为精，仁爱救人

## CONTENTS

INTRODUCTION .....	1
1. VASCULAR ENDOTHELIAL GROWTH FACTOR B.....	1
1.1 Biology, Tissue Distribution and Regulation of VEGF-B .....	1
1.1.1 The Biology of VEGF-B.....	1
1.1.2 Tissue Distribution.....	2
1.1.3 Regulation of VEGF-B.....	3
1.2 Role in Angiogenesis, Fatty Acid Uptake and Metabolism.....	4
1.2.1 Role of VEGF-B in Angiogenesis .....	4
1.2.2 Fatty Acid Uptake and Metabolism .....	5
2. ISLETS OF LANGERHANS AND LIPIDS .....	8
2.1 Islet Anatomy, Biology and Vasculature .....	8
2.1.1 The Anatomy and Biology of Islets of Langerhans .....	8
2.1.2 The Vasculature of the Islets of Langerhans .....	9
2.2 Lipid Transport and Pancreatic Islets.....	10
2.2.1 Endothelial Lipid Transport .....	10
2.2.2 Lipid Transport in Pancreatic Islets.....	12
2.2.3 Lipid Signaling in Pancreatic Islets .....	13
2.3 Lipotoxicity in Pancreatic Islets .....	15
2.3.1 ER Stress .....	15
2.3.2 Mitochondrial Dysfunction .....	16
2.3.3 Defective Autophagy.....	16
3. ISLET INTRAVITAL IMAGING.....	18
3.1 Development and History .....	18
3.2 Islet Intravital Imaging Technique and Application.....	19
4. ENDOTHELIAL Wnt/ $\beta$ -Catenin SIGNALING.....	21
4.1 Introduction to Wnt/ $\beta$ -Catenin Signaling .....	21
AIMS OF THE THESIS .....	23
RESULTS AND DISCUSSION.....	24
ACKNOWLEDGEMENT .....	32
REFERENCES.....	39

## LIST OF ABBREVIATIONS

ACE	Anterior chamber of the eye
CK1	Casein kinase
DAG	Diacylglycerol
DKD	Diabetic kidney disease
Dvl	Dishevelled
ER	Endoplasmic reticulum
ERR $\alpha$	Estrogen-related receptor $\alpha$
FA	Fatty acid
FABP	Fatty acid binding protein
FAT	Fatty acid translocase
FATP	Fatty acid transport protein
FFA	Free fatty acid
FFAR	Free fatty acid receptor
FZD	Frizzled
GPR	G-protein-coupled receptor
GSIS	Glucose stimulated insulin secretion
GSK3	Glycogen synthase kinase 3
LCFA	Long chain fatty acid
LDL	Low density lipoprotein
LRP	Low density lipoprotein receptor-related protein
MAG	Monoacylglycerol
NADH	Nicotinamide adenine dinucleotide, reduced form
NRF1	Nuclear respiratory factor 1
NRP 1/2	Neuropilin 1/2
PGC-1 $\alpha$	Peroxisome proliferator activated receptor $\gamma$ coactivator 1 $\alpha$
PPAR $\gamma$	Peroxisome proliferator activated receptor $\gamma$
ROS	Reactive oxygen species
SCFA	Short-chain fatty acid
TCF	T cell factor
TG	Triglyceride
UPR	Unfolded protein response
VEGF	Vascular endothelial growth factor
VEGFR	Vascular endothelial growth factor receptor
VPF	Vascular permeability factor



# INTRODUCTION

# I

## VASCULAR ENDOTHELIAL GROWTH FACTOR B


### 1.1 Biology, Tissue Distribution and Regulation of VEGF-B

#### 1.1.1 The Biology of VEGF-B

The vascular endothelial growth factor (VEGF), previously known as the vascular permeability factor (VPF), was first discovered by Senger et al. in 1983 through the study of tumor ascites fluids secreted by a number of tumor cell lines<sup>4</sup>. Later that year, Napoleone Ferrara successfully isolated VEGF and indicated its role in angiogenesis<sup>5</sup>. The VEGF family consists of six mammalian glycoprotein growth factors: VEGF-A (or VEGF), VEGF-B, VEGF-C, VEGF-D, VEGF-E and placenta growth factor (PlGF)<sup>6</sup>. They mainly bind to three different tyrosine kinase receptors that are expressed by vascular and lymphatic endothelial cells with an overlapping pattern: VEGFR1, VEGFR2 and VEGFR3<sup>7</sup>. VEGF is essential for vasculogenesis during embryonic development and also for angiogenesis under both physiological and pathological conditions<sup>6,7</sup>. In endothelial cells, the common semaphorin receptors Neuropilin-1 (NRP1) and NRP2 can also act as co-receptors for several of the VEGFs<sup>8</sup> (Figure 1). The VEGF ligands and their receptors mediate signaling in both autocrine and paracrine fashion<sup>9-11</sup>.

VEGF-B, a member of the VEGF family encoded by the *Vegfb* gene, was first discovered in 1996 and is highly conserved in mammalian species<sup>12,13</sup>. Structurally, the *Vegfb* gene contains seven exons and due to alternative splicing on splice acceptor sites in exon 6, the *Vegfb* gene

gives rise to two protein isoforms, VEGF-B<sub>167</sub> and VEGF-B<sub>186</sub><sup>12,14</sup>. Both VEGF-B isoforms are expressed in various tissues and binds specifically to VEGFR1 and NRP1 on endothelial cells<sup>15,16</sup>. VEGF-B<sub>167</sub> is the prevalent isoform and contains a highly basic heparin-binding carboxyl terminus, which promotes binding to cell surface heparan sulfate proteoglycans in the extracellular matrix<sup>17</sup>. In contrast, VEGF-B<sub>186</sub> contains a hydrophobic carboxyl terminus and is freely diffusible. Unlike VEGF-B<sub>167</sub>, proteolytic cleavage of the VEGF-B<sub>186</sub> carboxyl terminus is required before binding to the co-receptor NRP1<sup>15</sup>.



**Figure 1. Schematic diagram of VEGF ligands and receptors**

Partially overlapping binding pattern of VEGF ligands to their receptors. VEGF-A, VEGF-B and placental growth factor (PIGF) also bind the co-receptor NRP1. (Modified image from Hagberg *et. al*, *Physiology*, 2013')

### 1.1.2 Tissue Distribution

Under physiological conditions, VEGF-B is widely expressed in adult mouse and human tissue with abundant expression in tissues with high metabolic capacity, such as the myocardium, skeletal muscle, vascular smooth muscle, brown adipose tissue, kidney and brain, among which heart and skeletal muscle have the highest expression<sup>9,13,18-20</sup>. However, endothelial cells seem to lack VEGF-B expression<sup>20</sup>. VEGF-B<sub>167</sub>, which accounts for over 80% of total VEGF-B transcripts is expressed in most tissues in the mouse<sup>17</sup>. VEGF-B<sub>186</sub> isoform on the other hand, exhibits a much lower expression level and is expressed in a limited

number of organs<sup>17</sup>. Interestingly, contrary to the generally low expression of the VEGF-B<sub>186</sub> isoform in the normal healthy state, upregulated expression of VEGF-B<sub>186</sub> was found in both human and mouse tumor cell lines<sup>17</sup>.

With regard to VEGF-B specific expression in pancreas, VEGF-B has been shown to be expressed in pancreatic islets, including the  $\beta$ -cells, similar to other members of the VEGF family<sup>21,22</sup>. Single cell transcriptome studies on mouse and human islets also revealed that *Vegfb* is expressed in all types of endocrine cells and also in acinar cells<sup>23,24</sup>. In our recent study (listed as paper 1 in this thesis), we have additionally demonstrated that VEGF-B is expressed both in endocrine and exocrine tissues with similar abundance using RNA *In Situ* hybridization and that the expression level in pancreatic islets is approximately 10-fold lower compared to heart and skeletal muscle<sup>25</sup>.

Under pathological conditions, VEGF-B has been suggested to play a role in neoangiogenesis due to higher expression level in certain cancer types<sup>26</sup>. However, VEGF-B expression level remains constant through hyperplastic progression or angiogenic phase when crossed with pancreatic cancer inducing mouse model (RIP1-Tag2)<sup>21</sup>. VEGF-B transcripts were also found in various human cancers such as breast<sup>27</sup> and ovarian cancer<sup>28</sup>, adenocarcinoma<sup>29</sup>, and sarcoma, lymphoma and melanoma<sup>26</sup>. Furthermore, a decreased VEGF-B expression were reported in dilated cardiomyopathy and ischemic heart disease<sup>30</sup>.

### 1.1.3 Regulation of VEGF-B

Current knowledge regarding transcriptional regulation of VEGF-B expression is still limited. Of note, bioinformatic analysis revealed that *Vegfb* is tightly co-expressed with a cluster of nuclear-encoded mitochondrial genes, in particular NADH dehydrogenase 1a subcomplex 5 (*Ndufa5*) and cytochrome c (*Cyts*)<sup>19</sup>. Notably, these two genes are regulated by peroxisome proliferator activated receptor  $\gamma$  co-activator 1 $\alpha$  (PGC-1 $\alpha$ ), a master regulator of mitochondrial energy metabolism<sup>31</sup>. Therefore, these data hinted to the possibility that PGC-1 $\alpha$  might also regulate VEGF-B expression. A later study by Mehlem et al. indeed demonstrated that *Vegfb* expression is regulated by PGC-1 $\alpha$ <sup>32</sup>. In response to external stimuli such as exercise, fasting and cold, PGC-1 $\alpha$  is induced and activates several transcription factors, such as nuclear respiratory factor 1 (NRF1), PPAR $\gamma$  and estrogen-related receptor  $\alpha$  (ERR $\alpha$ ), to exert its effects<sup>32</sup>. PGC-1 $\alpha$  activation during fasting/exercise induces mitochondrial biogenesis, facilitates cellular fatty acid (FA) uptake for  $\beta$ -oxidation, and

promotes angiogenesis by increasing the expression of *Vegfa* and *Vegfb*<sup>32</sup>. Having VEGF-B expression under the control of PGC-1 $\alpha$  fits very well from an evolutionary point of view in order to ensure sufficient nutrient supply to tissues with high metabolic activities in order to maintain tissue energy homeostasis under challenging conditions.

Unlike other members of the VEGF family, VEGF-B expression was originally not thought to be induced by hypoxia due to lack of a hypoxia-response element sequence in the *Vegfb* promoter region<sup>33</sup>. However, a recent study demonstrated hypoxia-induced VEGF-B expression in mouse retina and that VEGF-B signaling via VEGFR1 plays a role in pathological retinal neovascularization<sup>34</sup>. Therefore, further studies are warranted to further validate the role of hypoxia in the regulation of VEGF-B expression.

## **1.2 Role in Angiogenesis, Fatty Acid Uptake and Metabolism**

### **1.2.1 Role of VEGF-B in Angiogenesis**


Attributable to the high sequence homology and overlapping receptor binding patterns to VEGF-A<sup>35,36</sup>, VEGF-B was originally thought to be an angiogenic factor, and VEGF-B was initially shown to stimulate endothelial cell growth *in vitro*<sup>13</sup>. However, unlike mouse models exhibiting VEGF-A or VEGF-C deficiency that are embryonically lethal with vascular defects, VEGF-B deficient mice are healthy and fertile<sup>37-40</sup>. Pathological angiogenesis in most organs studied are also not affected by VEGF-B deficiency<sup>41</sup>. Additional studies in deficient mice also have demonstrated that VEGF-B is neither required for vessel formation in proliferative retinopathy<sup>39</sup> nor for the blood vessel remodeling in pulmonary hypertension<sup>40</sup>.

Strategies to study VEGF-B's angiogenic potential by transgenic overexpression *in vivo* have mostly turned out negative<sup>42,43</sup>, in contrast to most other VEGF family members that exhibit either angiogenic or lymphangiogenic properties after transgenic overexpression, such as VEGF-A<sup>44-46</sup>, VEGF-C<sup>47</sup>, VEGF-D<sup>48</sup> and VEGF-E<sup>49</sup>. For instance, ectopic overexpression of VEGF-B under the Tie2 promoter in endothelial cells did not induce angiogenesis *in vivo*, and instead it was suggested that endothelial cell specific overexpression of VEGF-B potentiated postnatal vessel growth rather than initiation of vessel growth<sup>43</sup>. The use of adenoviral vectors for delivering VEGF-B into muscle and periadventitial tissue did not trigger vessel growth either<sup>50,51</sup>. In addition, although VEGF-B overexpression in the skin increased capillary diameter, the effect on blood vessel density was modest<sup>42</sup>.

In the eye, overexpression of VEGF-B by gene transfer has on the contrary been shown to promote pathological retinal and choroidal neovascularization in mice<sup>52</sup>. In addition, Robciuc et al. demonstrated a novel function of VEGF-B in adipose tissue, where gene transfer of the VEGF-B<sub>186</sub> isoform via an adeno-associated viral vector resulted in increased vascularity and perfusion of the adipose tissue, correlating with an improved insulin sensitivity and overall metabolic condition<sup>53</sup>. The positive effect of VEGF-B overexpression on angiogenesis was attributed to increased ligand occupation of VEGFR1, resulting in increased bioavailability of VEGF-A to VEGFR2 and activation of proangiogenic VEGF-A/VEGFR2 signaling<sup>53</sup>.

### 1.2.2 Fatty Acid Uptake and Metabolism

Hagberg et al. demonstrated that stimulation of endothelial cells with both VEGF-B isoforms induce expression of fatty acid transporter protein (FATP)-3 and 4 both at mRNA and protein level<sup>19</sup>. Stimulation of cultured endothelial cells with VEGF-B also increased the cellular uptake of long chain fatty acids (LCFAs), and that this process could be blocked by silencing the expression of FATP3 and FATP4 using siRNA<sup>19,54</sup>. Further *in vitro* studies revealed that treatment of cultured endothelial cells with VEGFR1 and NRP1 neutralizing antibodies blocked expression of FATP3, suggesting that both receptors were required for VEGF-B mediated FATP expression<sup>19</sup>. *In vivo* studies have also shown that overexpression of VEGF-B by adenoviral delivery is capable of inducing FATP expression<sup>19</sup>. Depletion of *Vegfb* in mice resulted in decreased expression of FATP3 and FATP4 in endothelial cells and decreased lipid accumulation in the heart, skeletal muscle and brown adipose tissue. Instead lipids were redirected to white adipose tissue<sup>19</sup>. In addition, both VEGFR1 and endothelial cell-specific NRP1 deficient mice exhibit defective FA uptake to peripheral tissues, indicating that both receptors are essential for VEGF-B mediated FATP expression *in vivo*<sup>19,20</sup>. Taken together, these data indicated that VEGF-B controls endothelial FA uptake through a paracrine regulation of vascular FATPs via VEGFR1 and NRP1 signaling (Figure 2).



**Figure 2. Schematic illustration of VEGF-B in trans-endothelial FA transport**

1: VEGF-B and mitochondrial proteins under the control of PGC1 are co-expressed in tissue cells to co-ordinate endothelial fatty acid transcytosis and tissue  $\beta$ -oxidation. 2: VEGF-B signals via VEGFR1 and NRP1 on the endothelial cells in a paracrine fashion. 3: Stimulation of endothelial cells with VEGF-B upregulates vascular FATPs expression and induces transport of fatty acids cross the endothelial layer into tissue cells. (Modified Image from Hagberg et. al, *Physiology*, 2013<sup>1</sup>)

A follow-up study revealed that VEGF-B deficient mice kept on high fat diet or crossed with type-2-diabetic *db/db* mice display increased insulin sensitivity and improved glucose homeostasis and plasma lipid profile. Furthermore, genetic deletion of *Vegfb* in diabetic mice showed increased cardiac glucose uptake, decreased ectopic lipid deposition in heart and skeletal muscle, and normalized blood glucose. This effect could also be replicated in *db/db* mice by pharmacological inhibition of VEGF-B signaling using neutralizing antibodies<sup>55</sup>. The *db/db* mice showed improved glucose tolerance,  $\beta$ -cell function and preserved pancreatic islet architecture. The overall lipid storage was reduced in heart, skeletal muscle and pancreas, however, not in liver<sup>55</sup>, suggesting that targeting VEGF-B signaling can limit peripheral lipid accumulation, hence restore tissue insulin sensitivity and  $\beta$ -cell function. Therefore, VEGF-B is essential for tissue lipid homeostasis and constitute a target for developing a novel treatment against metabolic syndrome. In line with these observations, other studies have also shown similar results. Kivelä et al. observed increased FATP4 expression in rat heart with VEGF-B overexpression and conversely decreased FATP4 expression in rat heart with VEGF-B deficiency<sup>56</sup>. In addition, a study by Karpanen et al. demonstrated that cardiac

specific overexpression of VEGF-B resulted in increased ceramide accumulation in the heart<sup>42</sup>.

In a subsequent study, Falkevall et al. further demonstrated that inhibiting VEGF-B mediated FA uptake could be therapeutically beneficial. Using multiple mouse models of diabetic kidney disease (DKD), the study showed that renal VEGF-B expression correlated with DKD disease severity and inhibition of VEGF-B either through genetic deletion or pharmacological inhibition could reduce renal lipotoxicity, prevent renal dysfunction and inhibit DKD development<sup>57</sup>.

In the study by Robciuc et al. they also observed that activation of VEGF-B/VEGFR1 pathway by gene transfer could induce weight loss and counteract metabolic complications in obese and insulin resistant mice, suggesting the therapeutic potential for targeting VEGF-B/VEGFR1 pathway for treating metabolic disease<sup>53</sup>. Gene expression data from mice and human revealed high expression of VEGF-B and its receptors in both white and brown adipose tissue<sup>58</sup>, suggesting that VEGF-B might play an important role in adipose tissue. In white adipose tissue of VEGF-A deficient mice, VEGF-B, FATPs 1-4 and brown adipose tissue specific genes were upregulated and the mice also have lower body weight when fed on high fat diet<sup>59</sup>. Another study has shown that cold stimulation downregulates VEGF-B and FATP3 expression in brown adipose tissue<sup>60</sup>. However, both studies were observatory, and the precise mechanism regarding this regulation is still unclear.

# II

## ISLETS OF LANGERHANS AND LIPIDS

### 2.1 Islet Anatomy, Biology and Vasculature

#### 2.1.1 The Anatomy and Biology of Islets of Langerhans

The pancreatic islets, also referred to as the islets of Langerhans, were first discovered by the German anatomist Paul Langerhans in 1869<sup>61</sup>. Dispersed inside the pancreas, the islets represent the endocrine part of the pancreas and consist of at least five different hormone producing endocrine cell types, including the glucagon producing  $\alpha$ -cells, insulin producing  $\beta$ -cells, pancreatic polypeptide producing  $\gamma$ -cell and somatostatin producing  $\delta$ -cells.  $\beta$ -cells account for the majority of the islet cells (70%) and the hormones produced by the different endocrine cells are directly secreted into the circulation to exert their respective functions.

The distribution of the different endocrine cells within the islet are species dependent<sup>62</sup>. In human islets the  $\beta$ -cells are intermingled with other endocrine cells, whereas  $\beta$ -cells in rodent islets are surrounded by a ring of  $\alpha$ -cells<sup>63</sup>. The islets are defined as micro-organs with a highly specialized and dense capillary vascular network. Islets are spread out over the entire pancreas and constitute around 4.5% of the total pancreas volume<sup>64</sup>. Despite a rather small volume, the islets receive around 10-15% of the pancreatic blood flow. The dense islet vascular network has an important physiological function as it facilitates the rapid exchange of glucose and



insulin between *e.g.*  $\beta$ -cells and blood, thereby supporting glucose sensing and insulin secretion<sup>62</sup>.

One of the important functions of pancreatic islet is to rapidly secrete insulin/glucagon into circulation in response to physiological signals.  $\beta$ -cells as the majority of the cell population in islet, their ability to secrete insulin is essential in maintaining systemic glucose homeostasis. In order to achieve this rapid response, insulin is released from stored granules via exocytosis through two phases<sup>65</sup>. The first phase characterized by rapid release of insulin secretory granule from readily insulin granule pool in response to elevated blood glucose, whereas second phase is characterized by a sustained slow release of newly formed granule<sup>66</sup>. Dysfunction in insulin release and failure to maintain glucose homeostasis have been implicated in the onset of both type 1 and 2 diabetes mellitus.

### **2.1.2 The Vasculature of the Islets of Langerhans**

The establishment of the specialized endothelium within the islet vasculature is a direct result of micro-environmental signals, especially proteins from the VEGF family. VEGF-A has been shown to play a crucial role in the formation of the islet vasculature<sup>67</sup>. During embryonal development, differentiating endocrine cells start to secrete ample amounts of VEGF-A, which in turn attracts endothelial cells to the site of the developing islets. The endothelial cells in turn secrete angiocrine factors, causing neighboring  $\beta$ -cells to proliferate, and this reciprocal interaction shapes the outline of the islet vascular bed resulting in a dense capillary network<sup>68,69</sup>. VEGF-A is not only essential for formation of the islet vasculature, but also for the development of endothelial pores, denoted fenestrae<sup>65</sup>. Compared to exocrine tissue, islet capillaries are highly permeable, which enables the rapid exchange of macromolecules and prompt insulin delivery into the vascular space, as well as for efficient interstitial flow in mature islets<sup>68,70</sup>. To further understand the role of VEGF-A in islet vascularization and fenestration, a number of studies using transgenic mouse models where VEGF-A was either specifically deleted in  $\beta$ -cells or in the entire pancreatic epithelium has been evaluated<sup>22,68,70</sup>. VEGF-A deficient islets exhibit a significantly reduced vascular network, capillary size and endothelial cell permeability; 10 times less fenestrated than control islets, these mice are also glucose intolerant<sup>67,70</sup>, presumably due to a severely reduced insulin output into the vascular system<sup>68</sup>. However, VEGF-A deficiency did not completely suppress blood vessel formation, indicating that other growth factors are also involved in islet vascularization<sup>70</sup>.

One of the important functions of the capillary network in islets is to support efficient secretion of different metabolic hormones directly into the blood stream<sup>71-73</sup>. Understanding the explicit route that the blood will follow within the islet is therefore of great importance in order to understand the communication between different islet cells. Currently, the precise organization of blood flow within the islet is still debatable. Nevertheless, three different models have been proposed<sup>74</sup>. One model using *in vivo* imaging in the mouse observed that the direction of blood is going from the  $\beta$ -cell rich islet core towards the non  $\beta$ -cells in the periphery<sup>73</sup>. However, the question still remains as to whether blood flow in human islets follows the same route, as the distribution of  $\alpha/\beta$ -cells differ between human and mouse islets. In addition, it appears that islet blood flow is regulated by a neuroendocrine loop and endothelial mediators such as nitric oxide<sup>75</sup>.

Dysfunction of the islet vasculature and islet inflammation has been implicated in the pathogenesis of type 2 diabetes. Several mechanisms have been proposed, for instance, increased islet capillary blood pressure and perfusion have been observed in short term hyperglycemia. This is likely due to a mechanism mediated by nitric oxide<sup>76,77</sup>. However, a sustained increase in capillary pressure and perfusion would lead to endothelial cell injury and thickening of the capillary wall, eventually resulting in decreased islet perfusion followed by impaired insulin secretion<sup>78,79</sup>. Endothelial cell activation and oxidative stress have been observed to lead to recruitment of monocytes to islets, which in turn cause islet dysfunction by secreting proapoptotic cytokines<sup>80</sup>. Studies in a Zucker type 2 diabetic rat model also suggested a pathogenic role of altered islet vasculature in the onset of type 2 diabetes<sup>81</sup>. In summary, development of an abnormal islet vasculature and islet vascular defects may also contribute to the development of type 2 diabetes.

## **2.2 Lipid Transport and Islets of Langerhans**

### **2.2.1 Endothelial Lipid Transport**

During fasting and feeding cycles, in order to sustain energy homeostasis, larger metabolites e.g. glucose and LCFAs must be transported from storage sites to different organs for utilization. Blood vessels are the major channels to enable this transportation. However, the mechanisms that regulate lipid transport is still poorly understood. Endothelial cells are

constantly exposed to circulating lipids and free fatty acids (FFAs) must penetrate the endothelial barrier in order to enable uptake in underlying tissue cells. Originally, a model of a flip-flopping mechanism was proposed, in which the hydrophobic FAs could diffuse freely through the plasma membrane<sup>82</sup>. However, evidences from several other studies indicated the existence of an active FA trans-endothelial transport system<sup>83</sup>. Moreover, another study observed that high concentration of FFAs could disturb the endothelial barrier, which facilitates the movement of low density lipoprotein (LDL) into arteries<sup>84</sup>. Hence, the detailed molecular mechanisms that govern endothelial FA uptake is still controversial to date and collective data indicate that both protein-mediated uptake and non-specific uptake may be involved<sup>85,86</sup>. Several proteins involved in endothelial FA uptake have been characterized, for example the scavenger receptor CD36 and the FATP protein family (FATPs 1-6)<sup>83</sup>.

CD36, also called fatty acid translocase (FAT), is expressed by monocytes, thrombocytes and endothelial cells in most metabolically active tissues, such as cardiac and skeletal muscle, and white and brown adipose tissue<sup>87</sup>. CD36 has an important function in recognizing and binding different lipids and lipid molecules<sup>88-91</sup>, and to facilitate cellular uptake of LCFAs<sup>92,93</sup>. Although the precise biological function of CD36 in endothelial cells is still under debate, several studies have demonstrated its role in endothelial lipid transport<sup>94-96</sup>. For example, the transcription factor PPAR $\gamma$  upregulates *Cd36* expression<sup>94,95</sup>. Absence of CD36 expression and reduced FA uptake to white adipose tissue have accordingly been observed in mice with endothelial specific *Pparg* deletion, which highly resemble the phenotype of *Cd36* knockout mice<sup>96</sup>. In another interesting study, cold exposure could potentially induce *Cd36* expression in brown adipose tissue to maintain thermogenesis by increasing cellular TAG accumulation<sup>60</sup>. A recent study also identified a FA binding site in CD36, partly overlapping with the binding site of oxidized LDL<sup>97</sup>.

The FATP family consists of 6 highly conserved mammalian multi-transmembrane proteins (FATPs 1-6)<sup>98-100</sup>. Initially, the first FATP was discovered by murine adipocyte cDNA expression cloning as a protein that facilitated LCFA uptake<sup>99</sup>, and subsequently additional FATP family members have been discovered and renamed<sup>101</sup>. FATP1 is mainly expressed in adipose tissue and heart<sup>19,99</sup>. FATP2 is expressed almost exclusively in the liver and kidney<sup>99,102</sup>. FATP3 seems specifically to be expressed within the vasculature<sup>55,103</sup>. FATP4 has a similar expression pattern as FATP1 but is also expressed in the intestine and endothelium. FATP4 is the only FATP that is expressed in the intestine<sup>19,104</sup>. Endothelial cells mainly express FATP3,

however, other FATPs are believed to be expressed at lower levels<sup>105</sup>. All FATPs have been demonstrated to increase cellular LCFAs uptake both *in vivo* and *in vitro*<sup>19,106,107</sup>. For instance, FATP3 and FATP4 co-expression in cultured endothelial cells lead to increased LCFAs uptake<sup>19</sup>. However, some studies argue that the effect *in vivo* is most likely due to the assistance from several other proteins which facilitates the uptake of LCFAs into endothelial cells<sup>108</sup>.

### 2.2.2 Lipid Transport in Pancreatic Islets

FAs play an important role in  $\beta$ -cell survival and secretory function. However, excessive exposure of  $\beta$ -cells to FAs will result in a series of negative effects, such as impaired insulin gene expression and secretion, and initiation of  $\beta$ -cell apoptosis, which all contribute to the development of  $\beta$ -cell failure<sup>109</sup>. Therefore, the comprehension of how FAs are being transported into  $\beta$ -cells lays the foundation for identifying future drug targets for  $\beta$ -cell lipotoxicity.

Depending on the chain length, FAs can cross plasma membrane into  $\beta$ -cell either by diffusion (e.g. short chain FAs (SCFAs)) or via cell surface transporter CD36 (LCFAs)<sup>110</sup>. Upon entry, fatty acyl-CoA synthetase activates incoming FAs to generate acyl-CoA, which then further undergoes  $\beta$ -oxidation for energy production<sup>111</sup>. Alternatively, acyl-CoA enters the glycerolipid/free fatty acid cycle to produce triglycerides (TGs), diacylglycerol (DAG), monoacylglycerol (MAG) and other lipid components<sup>112</sup>, or participates in the sphingolipid synthesis pathway to generate metabolites such as ceramides and sphingosine-1 phosphate<sup>110</sup>. In addition to the exogenous source of FAs,  $\beta$ -cell FAs can be derived from hydrolysis of intracellular TGs and phospholipids as lipoprotein lipase is also expressed in pancreatic islets<sup>113,114</sup>.

The process of transmembrane FA uptake into  $\beta$ -cells is mediated by plasma membrane fatty acid-binding proteins (FABP<sub>pm</sub>), which dissociates the FA from albumin at plasma membrane prior to binding to CD36<sup>115,116</sup>. Upon entering  $\beta$ -cell, cytoplasmic FABPs bind the FA and transport it to specific cellular compartments for further utilization<sup>117</sup>. Furthermore, this process has also been demonstrated in human islets and in an *in vitro* study using INS-1 cells<sup>118,119</sup>.

### 2.2.3 Lipid Signaling in Pancreatic Islets


In addition to the abovementioned transmembrane transport machinery, a group of G-protein coupled receptors on the  $\beta$ -cell surface, also referred to as FFA receptors (FFARs), have recently been discovered to function as signaling receptors for FAs, affecting a number of intracellular lipid signaling pathways. Currently four FFA receptors, FFAR1 (GPR40), FFAR2 (GPR43), FFAR3 (GPR41) and FFAR4 (GPR120) have been shown to be expressed in rodent and human  $\beta$ -cells<sup>120</sup>. Interestingly, the FFARs (1-4) exhibit chain length specificity (Figure 3). For instance, FFAR1 and FFAR4 specifically bind to medium and long-chain FAs, whereas FFAR2 and 3 only bind to short-chain FAs<sup>120</sup>.

Upon binding of the FA to its respective receptor, the complex exerts different effects on  $\beta$ -cell and systemic glucose homeostasis. For instance, binding of eicosatrienoic acid (C20:3) to FFAR1 (GPR40) facilitates glucose stimulated insulin secretion in  $\beta$ -cells<sup>121,122</sup>. However, opposing effects were observed in a number of *in vivo*, *in vitro* and human studies regarding the downstream effects of FFAR1. An *in vitro* study demonstrated protective effect of FFAR1 against lipotoxicity in rat insulinoma cells (INS-1)<sup>123</sup>, and overexpression of FFAR1 in  $\beta$ -cells improved glucose tolerance and glucose-stimulated insulin secretion (GSIS) *in vivo*<sup>124</sup>. In line with these observations, a single nucleotide polymorphism in the *FFAR1* locus in humans is associated with insulin secretory dysfunction<sup>125</sup>. Conflicting with these results, a study by Steneberg et al. showed that loss of FFAR1 in mice protected against obesity-induced hyperglycemia, glucose intolerance and hyperinsulinemia, whereas FFAR1 overexpression in  $\beta$ -cells resulted in  $\beta$ -cell dysfunction and diabetes<sup>126</sup>. Yet in another study, chronic exposure of palmitate to human islets resulted in decreased insulin content and secretion, and these effects could be prevented by a FFAR1 antagonist, ANT203<sup>127</sup>. The discrepancies in the different studies might be attributed to the difference in study design, such as duration of exposure, and species differences.

FFAR4 (GPR120) is the receptor for unsaturated FFAs and LCFAs and shows high binding affinity to docosahexaenoic acid (DHA) and  $\alpha$ -linoleic acid, which are both  $\omega$ -3 FAs<sup>128</sup>. Activation of FFAR4 increases glucagon-like peptide-1 (GLP-1) secretion from intestinal enteroendocrine L cells which in turn enhances insulin release from  $\beta$ -cells<sup>128</sup> and protects  $\beta$ -cell from apoptosis via activation of phosphoinositide 3-kinase-Akt (PI3K) and extracellular signal regulated kinase (ERK)<sup>129</sup>. FFAR4 knockout mice are furthermore glucose intolerant and an inactivating mutation in *FFAR4* has been linked with obesity in human<sup>130</sup>.

FFAR2 and 3 (GPR43, 41) are receptors for SCFAs, particularly FAs consisting of 3, 4 and 5 carbons. Activation of these receptors by SCFAs has been shown to inhibit insulin secretion via G protein coupling<sup>131</sup> and loss of both receptors in  $\beta$ -cells improve glucose tolerance and insulin release in high fat diet mice<sup>131</sup>. However, FFAR2 and 3 are essential for SCFA mediated GLP-1 secretion from intestinal enteroendocrine L cells, and consequently the global knockout of both receptors resulted in glucose intolerance *in vivo*<sup>132</sup>. Thus, the therapeutic strategy by targeting these receptors is debatable.

A number of nuclear receptors have also been demonstrated to function as receptors for FFAs. PPAR $\alpha$  can function as a receptor for LCFAs and upon activation, PPAR $\alpha$  increases FA oxidation and hence reduces lipid storage, protecting from lipotoxicity<sup>133</sup>. Additionally, farnesoid X receptor (FXR), a bile acid receptor, is also shown to regulate  $\beta$ -cell function in human islets<sup>134</sup>.



**Figure 3. Schematic diagram of fatty acid uptake into pancreatic  $\beta$ -cells and potential consequence**

Palmitate (PA) transmembrane transport by CD36 results in formation of increased ceramide and formation of lipid droplets (LD), which trigger cell stress responses, such as ER stress, mitochondrial dysfunction and defective autophagy. PA also activates FFA receptors resulting in impaired insulin secretion. (Modified image from Oh *et. al*, *Frontiers in Endocrinology*, 2018<sup>3</sup>)

## 2.3 Lipotoxicity in Pancreatic Islets

Obesity is one of the causes for the development of type 2 diabetes and commonly associated with insulin resistance<sup>135</sup>. In physiological condition,  $\beta$ -cells are very sensitive to rising circulating blood glucose and respond by releasing insulin to maintain a normal blood glucose level. Several nutrients can also trigger insulin release such as amino acids and fatty acids.

Depending on the concentration and duration, FAs can exert both positive and negative effects on  $\beta$ -cells, such as  $\beta$ -cell survival, insulin secretion and membrane composition<sup>136</sup>. Increased amounts of circulating FAs as a direct result of increased lipid accumulation and lipolysis in peripheral tissue is commonly associated with obesity<sup>137</sup>. Moreover, chronic exposure of FAs to islets severely suppresses the synthesis of proinsulin, impairs glucose stimulated insulin secretion from islets and also decreases insulin storage, leading to apoptosis and eventually results in  $\beta$ -cell failure<sup>136</sup>.

Dysfunction of the secretory capacity of  $\beta$ -cells and insufficient insulin production due to detrimental loss of  $\beta$ -cell mass eventually leads to the development of type 2 diabetes. Pancreatic islet lipotoxicity is considered to be one of the main causes for the onset and progression of this pathological process. Several likely mechanisms of lipotoxicity-induced  $\beta$ -cell dysfunction have been proposed over the years; in this chapter, endoplasmic reticulum (ER) stress, mitochondrial dysfunction and defective autophagy are mainly discussed.

### 2.3.1 ER Stress

In normal conditions, pancreatic  $\beta$ -cells depend on a highly active ER to satisfy the high secretory demands of insulin, as more than half of the total protein is synthesized in the ER<sup>138</sup>. Therefore,  $\beta$ -cell function is particularly vulnerable to ER stress, which subsequently would trigger unfolded protein response (UPR) on ER.

FA-induced ER stress is mainly mediated through interference with protein processing and trafficking<sup>139</sup> or dysregulation of  $\text{Ca}^{2+}$  efflux<sup>140</sup>. Disturbance of  $\text{Ca}^{2+}$  regulation in the ER results in incorrect protein folding, subsequently causing ER stress. Several studies have shown that long-term exposure of saturated FAs could inhibit ER vesicle trafficking in  $\beta$ -cells, confirming the theory that impaired protein trafficking could result in ER stress<sup>141</sup>. Palmitate (C16:0), the most common saturated FA, was recently shown to induce ER stress via depletion of ER  $\text{Ca}^{2+}$ <sup>142</sup>. And in the case of prolonged ER stress due to  $\beta$ -cell lipotoxicity,

failure of repairing incorrectly folded proteins may lead to  $\beta$ -cell apoptosis<sup>143</sup>. Hence, current findings indicate that lipotoxicity plays a unique role in ER stress mediated  $\beta$ -cell failure in the development of type 2 diabetes.

### 2.3.2 Mitochondrial Dysfunction

Dysfunction of mitochondria have been strongly correlated with impaired  $\beta$ -cell function. Under physiologically low glucose conditions,  $\beta$ -cells utilize FAs to generate ATP via  $\beta$ -oxidation to maintain their energy demands<sup>144</sup>. In obesity, increased availability of circulating FAs augment  $\beta$ -oxidation and oxidative phosphorylation in mitochondria, which result in excessive reactive oxygen species (ROS) production<sup>145</sup>. Pancreatic  $\beta$ -cells have a limited capacity for handling oxidative stress, therefore mitochondria, where the majority of ROS are produced, carries a higher risk for suffering from oxidative injury<sup>146</sup>. In normal conditions, only 0.1% of consumed oxygen generates ROS due to inefficient electron transport<sup>147</sup>. However, in diabetes and other pathological conditions, exacerbated ROS formation eventually results in oxidative stress and damaged mitochondria leading to  $\beta$ -cell failure<sup>146</sup>. Studies have shown that long term exposure of rat islets to saturated FAs induces  $\beta$ -cell apoptosis and this effect could be rescued by addition of antioxidants, suggesting that the toxic effect of FAs on  $\beta$ -cells could be mediated by oxidative stress<sup>148</sup>.

### 2.3.3 Defective Autophagy

Autophagy is an important cellular mechanism that allows the safe removal of dysfunctional cellular components, protein aggregates and pathogens. Recent research on the connection between intracellular lipids and autophagy in pancreatic islets indicated that autophagy also plays a role in the pathogenesis of type 2 diabetes. Under normal physiological conditions, autophagy is important for the maintenance of cellular quality control in  $\beta$ -cells, and this activity is indispensable for proper  $\beta$ -cell architecture and function<sup>149</sup>. Mice with deficient  $\beta$ -cell autophagy display with impaired glucose tolerance and insulin secretion, decreased  $\beta$ -cell proliferation, increased apoptosis and accumulation of protein aggregates<sup>150</sup>. The mice also exhibit accumulation of autophagosomes when evaluated in diabetic settings<sup>150</sup>. An *ex-vivo* study using non-diabetic human islets demonstrated build-up of autophagic vacuoles and increased apoptosis after exposure to saturated FAs<sup>151</sup>. In addition, several *in vitro* studies of INS-1E cells also observed increased amounts of autophagosomes when treated with saturated FAs, and this effect was also accompanied by impaired lysosome formation<sup>152</sup>.



Electron microscopic analysis of human islets from type 2 diabetic patients also revealed excessive autophagosomes which might be caused by impaired lysosome formation<sup>151</sup>. These data indicate that impaired  $\beta$ -cell autophagy may be more susceptible to lipotoxicity. However, further studies are needed to elucidate the specific role of autophagy in lipotoxicity induced  $\beta$ -cell failure.

# III

## ISLET INTRAVITAL IMAGING

### 3.1 Development and History


The pathophysiology of type 2 diabetes is owing to the development of  $\beta$ -cell dysfunction, reduction in  $\beta$ -cell mass and insulin resistance<sup>153-155</sup>. Major efforts have been investing into understanding the mechanisms of underlying causes of these processes within the research field. Notably, functional differences of pancreatic islets *in vitro* versus *in vivo* have been documented<sup>156,157</sup>. However, one major challenge is the lack of appropriate noninvasive animal models to study dynamic changes in islet mass and function during disease progression of *in vivo*. Approaches, such as *in vivo* optical monitoring of  $\beta$ -cell mass using MRI<sup>158</sup> lacks the spatial resolution required to study physiology at cellular level. Therefore, development and application of suitable and accessible model systems to enable *in situ* visualization and assessment of islet function *in vivo* is warranted.

The anterior chamber of the eye (ACE) provides an optically accessible site with a suitable innate environment in terms of oxygen supply, nerve innervation and immune system<sup>159</sup>. In 2008, Speier et al. developed the technique combining transplantation of pancreatic islets into the ACE and confocal/multiphoton microscopy to visualize and study islet function/behavior *in vivo*<sup>160</sup>. This minimal-invasive method makes it possible to visualize pancreatic islets under

different conditions in real-time<sup>160</sup>. Since its development, the islet intravital imaging setup has been widely used to address various questions within diabetes research.

### 3.2 Islet Intravital Imaging Technique and Application

The islet intravital imaging technique consists of two parts: First, freshly isolated or cultured pancreatic islets are transplanted into the ACE. This is followed by imaging of the engrafted islets at various timepoints post transplantation<sup>161</sup>. Depending on the research objectives, islets from donor mice harboring specific genetic modifications allowing for example direct visualization of a particular cell type (e.g. vessel specific labeling) can be transplanted to recipient mice<sup>159</sup>. To enable subsequent imaging of engrafted islets in the ACE, the anesthetized recipient mouse needs to be fixated on an imaging platform using a head holder, and then the eye is stabilized with an eyeball holder and oriented towards the microscope lens (Figure 4)<sup>161</sup>.



**Figure 4. Graphic illustration of noninvasive intravital imaging technique**

1. Isolation of tumorigenic RT2 islets. 2. Transplantation of isolated RT2 islets into the anterior chamber of the eye. 3. Weekly multiphoton/confocal intravital imaging of islet tumor progression. (Modified image from Balan et. al, *Scientific Reports*, 2019<sup>2</sup>)

The islet intravital imaging technique provides several benefits. Firstly, the minimal-invasive nature of this technique by using the eye as a natural window and therefore removing the anatomical barrier to enable *in vivo* imaging. Secondly, the high spatial imaging resolution that can be obtained by confocal and multiphoton imaging. Thirdly, the preservation of

morphological and functional characteristics of engrafted islets, which increase the relevance of the experimental observations both in physiological and pathological conditions<sup>157,159,160</sup>.

Since the development of this technique, substantial amount of knowledge has been accumulating regarding islet engraftment. Various studies have demonstrated that implanted islets rapidly integrate in the ACE environment and acquire adequate vascularization and innervation<sup>162-165</sup>. Furthermore, this technique is not limited only to islet transplantation, a recent study has also successfully demonstrated transplantation of pancreatic buds from 10.5-day mouse embryos for the purpose of studying pancreatic islet development<sup>166</sup>. Furthermore, a number of studies have been using this technique to investigate the *in vivo* dynamic profile of  $\beta$ -cell  $\text{Ca}^{2+}$  signaling, function and insulin resistance during the development of type 2 diabetes<sup>167-169</sup>.

In study III of this thesis, we subjected islets from a mouse model of pancreatic neuroendocrine tumor (Rip1Tag2, RT2 for short) to intravital imaging to study the development of tumor vasculature growth and dynamics with VEGF-B genetic manipulations. The RT2 mouse model is a well characterized pancreatic tumor model, in which simian virus 40 large T-antigen (Tag) oncogene is expressed under the rat insulin promoter. From birth, a multifocal stepwise  $\beta$ -cell carcinoma will develop in pancreatic islets<sup>170,171</sup>. Detailed description is described in paper III.

# IV

## ENDOTHELIAL Wnt/ $\beta$ -Catenin SIGNALING

### 4.1 Introduction to Wnt/ $\beta$ -Catenin Signaling

The term Wnt originated from the combination of Wingless and Int-1<sup>172</sup>. The Wnt signaling cascade is conducted either in paracrine or autocrine fashion and is highly conserved between species<sup>173,174</sup>. Currently three Wnt pathways have been identified: The canonical Wnt/ $\beta$ -catenin pathway, the noncanonical planar cell polarity pathway and the noncanonical Wnt/ $\text{Ca}^{2+}$  pathway. The canonical Wnt/ $\beta$ -catenin pathway is the most studied and regulates gene transcription<sup>175</sup>, the noncanonical planar cell polarity pathway regulates cytoskeletal organization and coordinate the polarization of the cell<sup>176</sup>, and the Wnt/ $\text{Ca}^{2+}$  regulates cellular calcium signaling and affects various intracellular signaling<sup>177,178</sup>. Wnt signaling pathways play a vital role in embryonic development, and is involved in a number of cellular processes such as cell fate specification, cell proliferation and migration, and genetic mutations affecting Wnt signaling have been implicated in cancer development<sup>175,179</sup>. In this chapter, only the canonical Wnt/ $\beta$ -catenin pathway will be discussed.

Activation of Wnt/ $\beta$ -catenin signaling requires the binding between Wnt and its receptor Frizzled (FZD) which leads to translocation of  $\beta$ -catenin to the nucleus and subsequent transcription of Wnt target genes<sup>174</sup>. In the absence of a Wnt ligand,  $\beta$ -catenin is targeted for proteolytic degradation, regulated by a destruction complex consisting of the tumor suppressor protein Axin, which functions as a scaffold for the complex, the tumor suppressor

protein APC, and two kinases, the casein kinase 1(CK1) and glycogen synthase kinase 3 (GSK3)<sup>180</sup>. CK1 and GSK3 sequentially phosphorylate  $\beta$ -catenin, and phosphorylated  $\beta$ -catenin is then recognized by the E3 ubiquitin ligase complex  $\beta$ -TrCP, which it ubiquitinates  $\beta$ -catenin and targets it subsequently targeted by proteasomal degradation, which resulting in rapid destruction of  $\beta$ -catenin<sup>181</sup>.

In the activated state, Wnt binds to its receptor FZD, and FZD then recruits the cytosolic effector protein Dishevelled (Dvl), which enables the binding of Axin to the plasma membrane. Axin together with other bound kinases phosphorylate low density lipoprotein receptor-related protein 5/6 (LRP5/6) in the plasma membrane, which provides further binding sites for Axin to form a stable Wnt-receptor-Dvl-Axin complex<sup>182</sup>. As a result, the destruction complex becomes inactivated, leading to accumulation of  $\beta$ -catenin in the cytosol and translocation to the nucleus where it interacts with DNA and T cell factor (TCF), resulting in transcription of Wnt target genes<sup>182,183</sup>.

Although the general theory of inner working of  $\beta$ -catenin destruction complex is widely accepted by the scientific community. New knowledge/evidence are emerging and continue to uncover additional intricate connections and regulations between each player. Along with the development of novel technologies, such as CRISPR-Cas genome editing system, it is likely to reveal more answers to many of the unsolved questions within the Wnt field in the foreseeable future.

## AIMS OF THE THESIS

The overall aim of this thesis is to investigate the role of VEGF-B in pancreatic  $\beta$ -cell physiology in the context of  $\beta$ -cell function, glucose homeostasis, islet lipid handling and angiogenesis. The specific aims are:

1. To investigate the effect of  $\beta$ -cell VEGF-B deficiency on pancreatic islet physiology, focusing on  $\beta$ -cell function and islet vascularization, and potential consequences on systemic glucose homeostasis and islet lipid content.
2. To study the impact of transgenic overexpression of the human VEGF-B<sub>167</sub> isoform in pancreatic islet  $\beta$ -cells on islet physiology and vascularization, and potential effects on systemic glucose and lipid homeostasis.
3. To explore the effect of human VEGF-B<sub>167</sub> overexpression on pancreatic islet tumor progression and tumor angiogenesis *in situ* using an in-house developed intravital high-resolution imaging platform.
4. To elucidate the role of endothelial Wnt/ $\beta$ -catenin signaling in central nervous system angiogenesis and potential crosstalk with VEGF signaling pathways.

## RESULTS AND DISCUSSION

### Study I. VEGF-B ablation in pancreatic $\beta$ -cells upregulates insulin expression without affecting glucose homeostasis or islet lipid uptake

Increased lipid uptake and handling can result in  $\beta$ -cell lipotoxicity, which contributes to  $\beta$ -cell dysfunction and development of type 2 diabetes mellitus. VEGF-B signaling in endothelial cells have been shown to increase endothelial fatty acid transcytosis via paracrine regulation of FATP-3 and -4, contributing to lipotoxicity in peripheral tissues such as heart and skeletal muscle. Systemic inhibition of VEGF-B results in improved metabolic performance in diabetic mouse models, correlating with improved insulin sensitivity and glucose tolerance<sup>55</sup>. However, the specific role of VEGF-B in islet  $\beta$ -cell physiology has yet to be examined. In this study, we generated and characterized a novel mouse model where VEGF-B in pancreatic  $\beta$ -cells has been selectively deleted using the Cre/LoxP system to investigate the tissue specific role of VEGF-B. We assessed whether ablation of VEGF-B in pancreatic  $\beta$ -cells would affect islet function, systemic glucose homeostasis and more importantly whether deficiency of VEGF-B would affect pancreatic islet lipid uptake, of relevance for  $\beta$ -cell lipotoxicity.

#### VEGF-B is ubiquitously expressed in pancreas

To address these questions, we started out by generating a mouse strain where *Vegfb* is selectively deleted in pancreatic  $\beta$ -cells. To achieve this, we crossed *Vegfb* floxed mice (*Vegfb*<sup>fl/fl</sup>) with RIP-Cre<sup>+/-</sup> mice, to generate *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> mice. To validate  $\beta$ -cell specific *Vegfb* deletion, RNA was isolated from *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> islets, and real-time quantitative PCR indicated 80% reduction of total *Vegfb* transcripts. Additionally, we also applied RNA *in situ* hybridization (RNAscope) to visualize the distribution of *Vegfb* transcripts. We observed that *Vegfb* is abundantly expressed in both the endocrine and exocrine parts of the pancreas in control mice, and that a visible reduction of *Vegfb* transcripts was confined to the islets in  $\beta$ -cell *Vegfb* deficient mice. Together these data indicated that *Vegfb* is abundantly expressed in pancreas and that the efficient reduction



of islet *Vegfb* transcripts in *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> mice points to that  $\beta$ -cells are the major source of islet VEGF-B.

### **Deficiency of *Vegfb* in pancreatic $\beta$ -cells results in increased insulin gene expression**

To evaluate the effect of *Vegfb* deficiency on islet gene expression, we performed real-time PCR on islet RNA retrieved from *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> and control mice. Interestingly, insulin gene *Ins2*, but not *Ins1*, was upregulated upon ablation of *Vegfb* in  $\beta$ -cells and conversely, we observed a trend of downregulation of islet glucagon gene expression (*Gcg*) when compared to the RIP-Cre control group but failed to reach significance when compared to the *Vegfb*<sup>fl/fl</sup> control. Furthermore, we also evaluated islet gene expression after 20-weeks of HFD feeding. However, islet insulin and glucagon expression levels remained similar across the groups.

Unlike other species, rodent insulin is encoded by two-gene system, *Ins1* and *Ins2* respectively<sup>184,185</sup>. *Ins1* is specifically to rodent, whereas *Ins2* has greater similarities to the *Ins* gene in other mammalian species. Insulin gene expression is tightly controlled by blood glucose via transcriptional regulation<sup>186</sup>, a number of transcriptional factors such as Pdx-1, PAX6, NueoD1 and MafA have been shown to regulate insulin expression in  $\beta$ -cell<sup>187-189</sup>. Our finding of *Vegfb* deficiency in  $\beta$ -cells only upregulates islet *Ins2* expression might suggest a potential link between *Vegfb* and transcriptional factors of *Ins2*. As PGC-1 $\alpha$  regulates VEGF-B expression<sup>32</sup>, and PGC-1 $\alpha$  has also been implicated in re-programming of Pdx-1 in adult  $\beta$ -cell<sup>190</sup>, hence it is plausible to speculate the possibility. Further studies are needed to elucidate aforementioned regulation machinery.

### **$\beta$ -cell specific *Vegfb* deficiency does not affect islet function or glucose homeostasis under chow and HFD diet condition**

To evaluate the effect of  $\beta$ -cell *Vegfb* deficiency on islet function, we examined glucose stimulated insulin secretion (GSIS) on isolated islets from all experimental groups. We found that  $\beta$ -cell *Vegfb* deficiency causes increased basal insulin release under low glucose conditions. However, this effect was diminished when stimulated with high glucose. Additionally, deficiency of *Vegfb* in  $\beta$ -cells did not alter total islet insulin content, or the pool of readily released insulin granules. When challenged by HFD, we did not observe any difference in GSIS, islet insulin content and KCl stimulated insulin secretion.

To investigate the effect of  $\beta$ -cell *Vegfb* deficiency on systemic glucose homeostasis, we measured a range of glucose related parameters, such as postprandial/fasting blood glucose, postprandial plasma insulin, glycated hemoglobin A<sub>1</sub>C, as well as performed glucose and insulin tolerance tests. Although we found significant changes in mice with  $\beta$ -cell *Vegfb* deficiency in postprandial/fasting blood glucose, postprandial plasma insulin level and insulin tolerance test when compared with RIP-Cre control groups, however, those parameters failed to reach statistical significance when compared to *Vegfb*<sup>fl/fl</sup> control mice. Similar set of tests were also performed on HFD mice, and consistent with previous observations, mice on HFD developed significant weight gain and mildly increased blood glucose, compared to age-matched mice on chow diet. However, all tested parameters related to glucose homeostasis showed insignificant difference in mice with  $\beta$ -cell *Vegfb* deficiency when compared to the control groups. Therefore, we concluded that deficiency of *Vegfb* in pancreatic  $\beta$ -cells does not affect islet function or overall glucose homeostasis under chow and HFD conditions, and increased insulin release under low glucose condition are likely attributed by cell autonomous effect and potentially conferred by upregulated islet *Ins2* gene expression.

### **Pancreatic islet VEGF-B expression is not affected by increased metabolic activity and $\beta$ -cell *Vegfb* deficiency does not affect lipid uptake in islets**

To understand the role of VEGF-B deficiency for islet lipid uptake and also the effect of  $\beta$ -cell *Vegfb* deficiency on downstream signaling of VEGF-B, we examined mice under both chow and HFD feeding conditions. Real-time PCR revealed that islet *Vegfb* transcripts were not increased after HFD treatment, suggesting that *Vegfb* expression is not regulated by increased metabolic activity in pancreatic islets. Plasma TG levels were similar between *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> mice and the controls groups both under chow and HFD. When comparing the TG level in mice fed on chow vs. HFD, increased plasma TG were found in all four genotypes. However, only RIP-Cre<sup>+/-</sup> and *Vegfb*<sup>fl/fl</sup>/RIP-Cre<sup>+/-</sup> groups reached a significant increase in circulating TG levels in HFD as compared to chow condition. To further examine the effect of  $\beta$ -cell *Vegfb* deficiency on islet lipid uptake, we measured islet ceramide content and *Plin2* gene expression, as both readouts represents possible fates of FA transportation into  $\beta$ -cells. Islet ceramide levels and *Plin2* gene expression were not affected by  $\beta$ -cell *Vegfb* deficiency under chow or HFD conditions. Additionally, *Cd36* and VEGF-B signaling downstream target genes encoding FATP3 and 4 were also unaffected.

Collectively, in this study we demonstrated that although *Vegfb* deficiency in islet  $\beta$ -cells correlated with upregulation of insulin gene expression, the effect on islet function, systemic glucose homeostasis or islet lipid uptake was negligible suggesting local paracrine VEGF-B signaling in pancreatic islets has limited therapeutic potential on overall glucose homeostasis and also as a target for reducing islet lipotoxicity.

## **Study II. Overexpression of VEGF-B in pancreatic $\beta$ -cells does not affect glucose homeostasis but increases plasma triglycerides**

Previous investigations have demonstrated a limited role for VEGF-B in angiogenesis<sup>2,21,53,54,56,191</sup>, but instead has an essential role for endothelial fatty acid transcytosis and lipid accumulation in tissues with high metabolic capacity been described<sup>55</sup>. However, the tissue specific role of VEGF-B signaling and whether local VEGF-B signaling would affect target organ function or systemic metabolic homeostasis remains unsettled. In this study, we have utilized a previously established mouse model where the human VEGF-B<sub>167</sub> isoform is transgenically overexpressed in  $\beta$ -cells, and we have assessed the effect under both chow and HFD conditions.

### **Transgenic overexpression of VEGF-B does not affect islet function or morphology**

We first validated the transgenic VEGF-B expression in pancreatic islets using real-time PCR. In comparison to wildtype control mice, islets from  $\beta$ -cell VEGF-B overexpressing (RIP1-VEGFB) mice showed similar amounts of mouse *Vegfb* transcripts, whereas islets from VEGF-B global knockout (*Vegfb*<sup>-/-</sup>) mice, as expected, had nearly no detectable mouse *Vegfb* expression. Furthermore, we observed that human *VEGFB* was about 40-fold higher expressed than mouse *Vegfb* in islets from RIP1-VEGFB mice, indicating an efficient transgenic overexpression of human *VEGFB*.

Next, we evaluated the islet function and morphology in wt and RIP1-VEGFB mice. Under both chow and HFD condition, GSIS revealed no difference in response to low and high glucose stimulation between wt and RIP1-VEGFB islets, and KCl stimulated insulin secretion also showed no difference. Taken together, these data suggested that  $\beta$ -cell VEGF-B overexpression does not affect islet secretory function. To further evaluate whether overexpression of VEGF-B affects pancreatic islet morphology, tissue sections from cryopreserved pancreata were subjected to immunofluorescence stainings with

insulin, glucagon and CD31 antibodies. Interestingly, we observed an increased vessel area (CD31 staining) in RIP1-VEGFB mice, which was in line with the observations from previous studies<sup>2,21</sup>. This effect could be explained by the VEGF-B/VEGFR1 decoy theory proposed by Robcius et. al<sup>53</sup>.

### **Systemic glucose homeostasis is not affected by VEGF-B overexpression in pancreatic $\beta$ -cells**

To address the question whether VEGF-B overexpression in pancreatic  $\beta$ -cell would affect systemic glucose homeostasis we measured a series of glucose parameters under both chow and HFD diet condition. We found that body weight, postprandial/fasting blood glucose, postprandial plasma insulin and glucose and insulin tolerance tests were all similar between wt and RIP1-VEGFB mice, suggesting that pancreatic  $\beta$ -cell overexpression of VEGF-B does not affect systemic glucose homeostasis.

### **Transgenic overexpression of VEGF-B in pancreatic $\beta$ -cells increases plasma triglycerides**

We unexpectedly found a significantly increased plasma TG level in RIP1-VEGFB mice. This phenotype was evident in both chow and HFD feeding condition. Interestingly, the increased TG level only existed in male mice, as plasma TG in female RIP1-VEGFB mice were of similar level to their wt control (data not shown). Currently, the particular mechanism of transgenic overexpression of VEGF-B in  $\beta$ -cells causing increased plasma TG is unknown. However, there are several possible angles that might help to explain this phenotype. As plasma TG is represented by both chylomicron and very low-density lipoprotein (VLDL), it is reasonable to speculate that pancreatic overexpression of VEGF-B can influence the formation of chylomicrons or VLDL, or alternatively, affecting activation of lipolysis. Further studies are warranted to elucidate this effect.

In summary, in this study we found that transgenic overexpression of VEGF-B in  $\beta$ -cells does not affect islet function and systemic glucose homeostasis. However, locally expressed VEGF-B in pancreatic  $\beta$ -cells causes increased vessel area and increased plasma TG level, a phenotype warranting further studies.

## **Study III. Noninvasive intravital high-resolution imaging of pancreatic neuroendocrine tumours**

In study III, we explored the role of the VEGF-B in tumor angiogenesis by employing a novel intravital 3D imaging platform developed in-house on islets from the pancreatic neuroendocrine tumor model Rip1Tag2, hereafter denoted RT2. The core principle of this technique is to transplant islets from mice with different genetic background (normal, fluorescent, tumorigenic) into the anterior chamber of the eye (ACE) of a recipient mouse, and using the eye as a natural window to image the engrafted islets with focus on vascularization, overall growth of tumor islets, the effect of VEGF-B on tumor angiogenesis and testing pharmacological inhibition with a topical approach and its effects on tumor angiogenesis.

### **Pancreatic neuroendocrine tumor characterization**

We first characterized RT2 tumor islet growth with ACE intravital imaging. Compared to normal islets, RT2 tumor islets grew rapidly, and were characterized by a tortuous vasculature. In addition, we observed that capillaries in RT2 tumor islets have irregular shapes, wider diameter and irregular blood flow. To further monitor tumor angiogenesis during cancer progression, we cross-bred RT2 mice with a number of different fluorescent reporter mice (EYFP, tdTomato) and injected fluorescence dyes to better visualize tumor angiogenesis and tumor cell behavior.

We further developed a tool for 3D quantification of ACE engrafted tumor progression, capable of calculating the tumor volume by 3D surface projection rendering, as well as by digital skeletonization of the islet vasculature assessment and quantify the number of branch/end points, segment diameter, length and number.

### **Sunitinib treatment resulted in tumor regression and reduced tumor growth**

Next, we validated the feasibility of pharmacological treatment of RT2 tumor in our ACE platform, with the objective of evaluating the effect of drug treatment of RT2 tumors in real-time. Two weeks after transplantation, pre-oncogenic RT2 islets were subjected to Sunitinib treatment for another two weeks and we observed strong tumor vessel regression in response to Sunitinib. Further quantification of Sunitinib treated RT2 tumor islets showed significantly decreased vascular volume, length and branch points. Control treated RT2 tumor islets continued to grow till the end of study, whereas Sunitinib treatment halted RT2 tumor growth. Taken together, these observations demonstrated the potential of the ACE platform for revealing novel information during the course of cancer

treatment, which could be beneficial for developing new strategies for future cancer treatment.

### **Overexpression of VEGF-B reduces tumor growth but increases tumor angiogenesis**

Based on previous studies, the role of VEGF-B in angiogenesis seem to be restricted to pathological conditions<sup>26</sup>. An earlier study found that overexpression of VEGF-B in RT2 tumor islets resulted in reduced tumor volume<sup>21</sup>. Endpoint histological analysis of VEGF-B overexpressing RT2 islets revealed furthermore no major vascular effects besides increased capillary diameter<sup>21</sup>. To gain further understanding of the role of VEGF-B in tumor angiogenesis, we crossed RIP1-VEGFB mice (mice from Paper II) with RT2 mice to generate double transgenic RT2-VEGFB mice.

Four weeks after ACE transplantation, we observed a reduction in tumor expansion in VEGF-B overexpressing RT2 islets (RT2-VEGFB). The tumors exhibited on the other hand an augmented angiogenic response. Further 3D quantification analysis indicated that RT2-VEGFB tumors display with increased vascular volume, length and number of branching points. However, vessel diameter was similar in RT2-VEGFB islets when compared to control RT2 islets. Interestingly, 3D quantification analysis of non-tumorigenic RIP1-VEGFB islets implanted into the ACE also showed increased vascular volume, number of branching points, as well as wider diameter, compared to control islets.

In summary, in this study we further developed the ACE platform and complemented with a more detailed quantification protocol, demonstrating the feasible application of the ACE platform in evaluating the efficacy of pharmacological treatments on tumor islets. More importantly, we showed that overexpression of VEGF-B suppresses pancreatic tumor islet growth and that this was accompanied by increased tumor angiogenesis.

### **Study IV. Endothelial $\beta$ -catenin signaling supports postnatal brain and retinal angiogenesis by promoting sprouting, tip cell formation, and VEGFR2 expression**

In this study, we explored the contribution of endothelial Wnt/ $\beta$ -catenin signaling in postnatal central nervous system (CNS) angiogenesis. Activation of endothelial  $\beta$ -catenin signaling has an essential role in vascularization of brain and retina and mutations in the

endothelial  $\beta$ -catenin pathway has been linked to impaired vascularization, blood-brain barrier dysfunction and a number of CNS diseases<sup>192</sup>.

To visualize active  $\beta$ -catenin signaling in postnatal cerebral vasculature, we utilized a reporter mouse line where  $\beta$ -catenin transcriptional activity induces GFP expression and we observed that endothelial  $\beta$ -catenin signaling was very active throughout postnatal cerebral development. To better understand the role of Wnt/ $\beta$ -catenin in CNS angiogenesis, we used inducible genetically modified mice modeling activated (gain-of-function) or inactivated (loss-of-function) endothelial  $\beta$ -catenin signaling. Endothelial  $\beta$ -catenin loss-of-function in the postnatal stage (day 6) resulted in retinal hypovascularization with deficient endothelial cell proliferation and sprouting, in agreement with previous observations<sup>193,194</sup>. Conversely, using inducible  $\beta$ -catenin gain-of-function mice, we further demonstrated that endothelial  $\beta$ -catenin signaling regulates tip cell selection during sprouting angiogenesis in the retina.

Additionally, endothelial  $\beta$ -catenin signaling also actively promote blood-brain barrier formation and sprouting angiogenesis, as judged by the reduced expression of markers for these processes in  $\beta$ -catenin loss-of-function mice. Some of the effects may be linked to defective VEGF-A signaling as the expression of both VEGFR2 and -3 was diminished in the CNS microvasculature. Interestingly, we also found that impaired endothelial  $\beta$ -catenin signaling prevented pathological hypersprouting when Notch signaling was inhibited, presumably related to failure of upregulating VEGF-A/VEGFR2 signaling.

In summary, in this study we demonstrated that endothelial  $\beta$ -catenin is crucial for angiogenesis under low Notch signaling. We further propose that impaired endothelial  $\beta$ -catenin signaling result in reduced retinal and cerebral angiogenesis due to reduced VEGFR2 expression in the CNS microvasculature, demonstrating a crosstalk between  $\beta$ -catenin and the VEGF/VEGFR2 signaling pathway.

.

## ACKNOWLEDGEMENT

To learn new knowledge, to be able to comprehend and to exercise it to its full potential and ultimately to produce new knowledge reflect the essence and the journey of becoming a scientist. Over the years, I have been fortunate enough to learn from some of the best and brilliant people in the field, whom have been working tirelessly to generate the knowledge that I have benefited during my time as a doctoral student and also to build this thesis upon. There is no doubt that I will be forever grateful for the knowledge that I have learned during my time at Karolinska Institutet, but unforgettably the people that I have the privilege to meet and work with both professionally and personally deserve my unreserved gratitude.

Firstly, I would like to thank my supervisor **Daniel**, thank you for accepting me as a member of your group, and thank you for your constant support (both financially and mentally) throughout my PhD period and also for providing me any assistance whenever I needed. And for always being there for me, no matter a small chat or causal scientific discussion.

To my co-supervisor **Ingrid**, I cannot thank you enough for all your help during the editorial stage of my papers and thesis. Your thoroughness and meticulousness on scientific writing and reasoning always impressed many of us, and I truly enjoyed the causal scientific discussion we had over the past several months, and I am grateful for all the assistance and guidance you have provided.

To my co-supervisor **Ulf**, the best/coolest professor in town. Thank you for your unwavering support throughout these years, for sharing your vast knowledge, and for all the fun afterwork scientific discussions with wide spectrum of topics. More importantly, for your caring approach, understanding and guidance through my years as a PhD student, and I have learned a great deal from you.

To **Lars Muhl**, thank you for all your technical assistance and critical reading of my paper. Your input helped me solved many technical problems that I encountered, and your scientific attitude is always inspiring. To **Aga**, thank you for taking care of me when I first started, for your friendship, and the laughter. Thank you for always calming my nerve down when things did not work, and I hope by now I have won the competition with your avocado plant and make you proud! (well about the plant... it got lost during the move to Biomedicum). To



**Dominic**, the best physicist I have ever met. Although your time in the lab was short, thanks a million for all the afterwork drinks, and your help with microscope and for your firm belief of the posh within my dressing style (although I disagree most of your claims), and for always being a good friend. To **Lady Mirela**, thank you for your help with all the islets picking during past several years, for sharing all funny jokes and laughter and wish you all the best for your study at U town. Also, to **Nina** and **Marta**, for your experimental help during my PhD study. To my summer students **Jiarui Mi** and **William Lindström**, thank you for your staining work!

Many colleagues in Eriksson's group and division of vascular biology have provided invaluable support to make this thesis possible. I would like to thank **Annelie**, for your rigid scientific input on my projects and critical advices, for your joyful positive laughter that made me to believe that I can conquer every challenging experiment. To **Sofia**, the best singer in the group, thank you for your help with all the mice experiment and your cheerful small talks and being the best problem-solver, thank you for the critical saves on many of my experiments. To **Annika**, for all the scientific input and advice on my projects and wish you all the best in your medical study. To **Erika**, thank you for taking care all of us, your work has made this lab work seamlessly! To **Björn**, for your daily cheerful laughter. To **Benjamin**, thank you for sharing many fun moments since the beginning of my doctoral study, I have learned the art of loyalty from your never-changing, consistent, stubborn and "I don't care how others think" cottage cheese only lunch-eating routine. **David Kaluza** for all funny jokes always make my day! Also, to current and former vascular biology members: **Isodle**, **Mannuel**, **Hanna**, **Karin**, **Aranzazu**, **Jongwook**, **Christine**, **Jonas**, **Maryam**, **Natalie**.

To **Linda Fredriksson**, thanks for all your advices and fun discussions during my group presentations. **Christina**, for sharing the PhD journey (although you finished way earlier than me). **Jil**, for being my desk neighbor in the office and your impressive organizational skills. And to **Lars Jakobsson**, thank you for all the scientific input on my projects and for guiding me through lipid-lab teaching. To **Maria** and **Antonio**, thank you for being awesome office buddies, and wish you all the best on your PhD journey. And to the people at 6D quarter, **Polina** and **Dimmitry**, thank you for organizing to all the nice parties!

And this thesis would not be possible without the amazing help from our wonderful KMB technician **Sara**! Thanks a million for your patience, and your preciseness when carrying out difficult tasks, and for always being there whenever I need your help! You are the true hero behind all these studies!

During the years, I have the privilege to meet some of amazing administrative personnel who make this journey as smooth as possible. Specially thanks to **Gizella**, for all your help with all the travel bookings and some rather challenging administrative requests and thank you for taking care all of us. As a PhD student in our department, we are lucky to have the two of the amazing doctoral study administrators, **Alessandra** and **Victoria** thank you for your caring personalities, and for always making sure we are on track to reach the milestones that set in our studies.

I also would like to thank **Prof. Monica Nistér**, for your acceptance and guidance during my time at your group. **Prof. Jian Zhao**, thank you for taking care of me during my master years and the solid training you provided. **Shaobo Jin** for welcoming to your home for dinner. **Mikael**, thank you for your help and advice. And to the former and current Nistér lab members, **Rong Yu, Tong Liu, Karl, Kaveh, Salah, Xiaobing, Inga, Tamador, Joseph, Anna, Elisa**, thank you all for your companies!

Over the years, I also have the honor to work and travel with some of amazing staffs at KI central administration. To **Emma Hägg**, thank you for the time working as student ambassador with you. To **Maissa**, thanks for inviting me on the Saudi trip! **Kristoffer**, thank you for being the awesome travel buddy and the fun afterworks! Wish you all the best with the new chapter into the fatherhood! To the dream team of China trip, **Prof. Lars-Arne Haldosen**, it was great to meet you on the trip, thank you for all the nice discussions about life and science, and your advices on going through my PhD, and of course, thank you for your signature on my degree certificate! To **Jenny**, it was great pleasure traveling and working with you over the years, and congratulations for your new life as a mother. To **Martin**, thanks for being the awesome travel buddy, thanks a million for all the trips we have been all these years and congratulations on your new life with Jenny! To **Katarina**, it was a pleasure to work with you on the recruitment trip and the fun cardiac development discussion over the plane! To **Monika**, thanks for inviting me on last year trip, and it was a pleasure to meet you and Stefan! **Prof. Robert Harris**, thank you for your time as our doctoral study director and your inspiration on what a student should achieve during your doctoral study, and it was a pleasure to work with you on last year trip. To **Prof. Nailin Li**, thank you for your advices and guidance over the years, and being a role model for many of us. And to **Maria**, thanks for all the work trips we have been on together and lunch chat over the topic of “finding the love of my life”, and more importantly thank you for encouraging me to get better with my Swedish, jag hoppas att jag en dag kommer att göra dig stolt över mig!

To my awesome friends during my master year: **Rasmus**, thank you for being a great friend and helping me settle in when I first moved to Stockholm. **Jie Ji, Bojing Liu, Jimmy Zhang**. To my friends at Biomedicum, **Xiao Tang, Chang Liu, Xiaoyuan Ren, Qing Cheng**.

To the Bobolinos crews: **Peter and Eilis, Will and Ella, Georgia and Gino, Olivia and Antonio, Christine and Robert, Carin and Alan, Marie** thanks for all the fun dinner parties, the fun weddings and babies' showers!

To the long island ice-tea crew: the brilliant artist and singer **Viktor, Alex and Moa, Johan and Malin, Manda**, thanks for your company during the sunny Malta trip and journey of finding the best long island ice-tea throughout the country.

To the magic **Mike Aron**, the awesome lipid lab teaching buddy for the past 3 years, thanks for always cheering me up at the end of exhausting day, and good luck with your doctoral study!

To **Hong**, thank you for listening and sharing when things didn't work out. And for all your help over the years, and good luck with your work! To **Ran Yang**, for organizing a great trip and all the fun dinner parties. **Xiong Yan and Yi Jin**, thank you for all the fun dinner parties and trips, thanks a million for listening and sharing my burdens when the days weren't so bright and for always be there for me. To **Yixin Wang**, my long-time loyal buddy, thank you for your great patience, your positivity and all the fun lunches. Thank you for making my days shining with your never-ending compliments and jokes, for always helping me whatever and whenever I need it. And congratulations on your new chapter in life!

To **Gioele**, the smartest Italian I have ever known. For the epic Japan trip, the memorable Akihabara afternoon, the Sunday ramen ritual and the rooftop bars about differential equations and seeing into the future. Thanks for being an awesome friend and for your constant inspiration for succeeding in science. Wish you all the best in your new career as a PI! To **Felix**, thanks for all the discussions of medical cases, the fun night out, and the elegant culinary adventures. Your passion towards medicine and your discipline will surely make you a great doctor and good luck with your new residency life!

To **Ragnar**, thanks for your nearly a decade friendship! For all the fun night outs, tasting of different cocktails, the late-night chicken wings tradition (or chilly cheese), and for always believing and following my motto of "Life is an adventure". Thanks a million for the journeys that we have been on together and I am looking forward to the next one!

To **Anna**, thank you for your friendship, the fun trips and always welcoming me at your home. I am happy that I have successfully persuaded you to join the KI track, and I look forward to our next journey (also next babysitting my lovely nephews)! To **Erika**, thank you for always remembering and celebrating my birthday, for always being there either a milestone in my life or the difficult times in all these years. I am happy we are going to defend our thesis only a few days apart, wish you best of luck for your defense and I am sure you will shine on that day! Also, to my lovely nephews: **Hugo, Timothy and Alfie**, I am proud to be your uncle and happy to see you growing bigger and bigger every day!

To **Peter Saliba Gustafsson**, for all these years, I am fortunate enough to have you as my friend. Thanks a million for the afterwork drinks, the Bagpiper's Inn, the brotherhood bonded during the missions of Call of Duty. Thanks for always sharing the burden when things did not work during my PhD, for being there in the time of sickness, and for encouraging and cheering me up and celebrating with me when I reached the milestone. Thanks for all the fun scientific discussions (although your wife thinks otherwise) and wish you all the best in Stanford, and I look forward to our next reunion!

To **O-J**, captain of the team, thank you for sharing every important moment during my time at Karolinska, and for your great friendship. You have always inspiring me with your novel ideas, your laser like focus and your self-discipline. For many people who know you would agree that you are a well accomplished researcher, but for me personally you are an exceptional friend. Thanks a million for always being there for me all these years. We attended our master graduation together and I am proud that we are going to do it again for our PhD!

借此毕业之时，感谢曾经教导过我的附属一院的老师们。感谢导师任延波教授多年来的悉心指导，感谢张彧主任为我提供临床学习的机会，感谢曾带教过我的晏辉老师，高冬娜老师，张雷老师。感谢心内科的刘莹教授，刘金秋教授，感谢心功能室的潘晓杰老师，曲丹丹老师，常青老师，王嘉慧老师，感谢脊椎外科的王宏教授。感谢老师们的悉心教导才有了我今天的进步。

感谢我在祖国家乡的兄弟连，悠悠、老猫、老大、三儿、潘书记、崔教授、DC 感谢大家二十年来不离不弃的友谊和多年来对我的支持！

感谢姑姑和叔叔、爷爷和奶奶、爸爸和妈妈从小到大无微不至的照顾和关爱，一如既往的支持和鼓励。

To my awesome brother **Ning Ning**, I am ridiculously proud to be able to grow up and with you and seeing you become a smart, responsible and fine young man that you are today. Thank you for sharing all the great memories growing up and I look forward to our next trip together!

Lastly, to my incredibly brilliant **Mom**, for always being a role model throughout my life, for your patience, understanding, encouragement, compassion and support. For being my loudest supporter, for holding me up when I fell, and for always loving me and believing in me.



## REFERENCES

- 1 Hagberg, C., Mehlem, A., Falkevall, A., Muhl, L. & Eriksson, U. Endothelial fatty acid transport: role of vascular endothelial growth factor B. *Physiology (Bethesda)* **28**, 125-134, doi:10.1152/physiol.00042.2012 (2013).
- 2 Balan, M. *et al.* Noninvasive intravital high-resolution imaging of pancreatic neuroendocrine tumours. *Sci Rep* **9**, 14636, doi:10.1038/s41598-019-51093-0 (2019).
- 3 Oh, Y. S., Bae, G. D., Baek, D. J., Park, E. Y. & Jun, H. S. Fatty Acid-Induced Lipotoxicity in Pancreatic Beta-Cells During Development of Type 2 Diabetes. *Front Endocrinol (Lausanne)* **9**, 384, doi:10.3389/fendo.2018.00384 (2018).
- 4 Senger, D. R. *et al.* Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* **219**, 983-985 (1983).
- 5 Leung, D. W., Cachianes, G., Kuang, W. J., Goeddel, D. V. & Ferrara, N. Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* **246**, 1306-1309, doi:10.1126/science.2479986 (1989).
- 6 Olsson, A. K., Dimberg, A., Kreuger, J. & Claesson-Welsh, L. VEGF receptor signalling - in control of vascular function. *Nat Rev Mol Cell Biol* **7**, 359-371, doi:10.1038/nrm1911 (2006).
- 7 Stuttfeld, E. & Ballmer-Hofer, K. Structure and function of VEGF receptors. *IUBMB Life* **61**, 915-922, doi:10.1002/iub.234 (2009).
- 8 Pellet-Many, C., Frankel, P., Jia, H. & Zachary, I. Neuropilins: structure, function and role in disease. *The Biochemical journal* **411**, 211-226, doi:10.1042/BJ20071639 (2008).
- 9 Aase, K. *et al.* Localization of VEGF-B in the mouse embryo suggests a paracrine role of the growth factor in the developing vasculature. *Dev Dyn* **215**, 12-25, doi:10.1002/(SICI)1097-0177(199905)215:1<12::AID-DVDY3>3.0.CO;2-N (1999).
- 10 Breier, G., Clauss, M. & Risau, W. Coordinate expression of vascular endothelial growth factor receptor-1 (flt-1) and its ligand suggests a paracrine regulation of murine vascular development. *Dev Dyn* **204**, 228-239, doi:10.1002/aja.1002040303 (1995).
- 11 Lee, S. *et al.* Autocrine VEGF signaling is required for vascular homeostasis. *Cell* **130**, 691-703, doi:10.1016/j.cell.2007.06.054 (2007).
- 12 Grimmond, S. *et al.* Cloning and characterization of a novel human gene related to vascular endothelial growth factor. *Genome Res* **6**, 124-131 (1996).
- 13 Olofsson, B. *et al.* Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 2576-2581 (1996).
- 14 Olofsson, B. *et al.* Genomic organization of the mouse and human genes for vascular endothelial growth factor B (VEGF-B) and characterization of a second splice isoform. *J Biol Chem* **271**, 19310-19317, doi:10.1074/jbc.271.32.19310 (1996).
- 15 Makinen, T. *et al.* Differential binding of vascular endothelial growth factor B splice and proteolytic isoforms to neuropilin-1. *J Biol Chem* **274**, 21217-21222, doi:10.1074/jbc.274.30.21217 (1999).
- 16 Olofsson, B. *et al.* Vascular endothelial growth factor B (VEGF-B) binds to VEGF receptor-1 and regulates plasminogen activator activity in endothelial cells. *Proc Natl Acad Sci U S A* **95**, 11709-11714, doi:10.1073/pnas.95.20.11709 (1998).
- 17 Li, X., Aase, K., Li, H., von Euler, G. & Eriksson, U. Isoform-specific expression of VEGF-B in normal tissues and tumors. *Growth Factors* **19**, 49-59, doi:10.3109/08977190109001075 (2001).
- 18 Lagercrantz, J. *et al.* A comparative study of the expression patterns for vegf, vegf-b/vrf and vegf-c in the developing and adult mouse. *Biochimica et biophysica acta* **1398**, 157-163 (1998).
- 19 Hagberg, C. E. *et al.* Vascular endothelial growth factor B controls endothelial fatty acid uptake. *Nature* **464**, 917-921, doi:10.1038/nature08945 (2010).
- 20 Muhl, L. *et al.* Expression of vascular endothelial growth factor (VEGF)-B and its receptor (VEGFR1) in murine heart, lung and kidney. *Cell Tissue Res* **365**, 51-63, doi:10.1007/s00441-016-2377-y (2016).

- 21 Albrecht, I. *et al.* Suppressive effects of vascular endothelial growth factor-B on tumor growth in a mouse model of pancreatic neuroendocrine tumorigenesis. *PLoS One* **5**, e14109, doi:10.1371/journal.pone.0014109 (2010).
- 22 Inoue, M., Hager, J. H., Ferrara, N., Gerber, H. P. & Hanahan, D. VEGF-A has a critical, nonredundant role in angiogenic switching and pancreatic beta cell carcinogenesis. *Cancer Cell* **1**, 193-202, doi:10.1016/s1535-6108(02)00031-4 (2002).
- 23 Segerstolpe, A. *et al.* Single-Cell Transcriptome Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes. *Cell Metab* **24**, 593-607, doi:10.1016/j.cmet.2016.08.020 (2016).
- 24 Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population Structure. *Cell Syst* **3**, 346-360 e344, doi:10.1016/j.cels.2016.08.011 (2016).
- 25 Ning, F. C. *et al.* VEGF-B ablation in pancreatic beta-cells upregulates insulin expression without affecting glucose homeostasis or islet lipid uptake. *Sci Rep* **10**, 923, doi:10.1038/s41598-020-57599-2 (2020).
- 26 Salven, P. *et al.* Vascular endothelial growth factors VEGF-B and VEGF-C are expressed in human tumors. *Am J Pathol* **153**, 103-108, doi:10.1016/S0002-9440(10)65550-2 (1998).
- 27 Ruohola, J. K. *et al.* Vascular endothelial growth factors are differentially regulated by steroid hormones and antiestrogens in breast cancer cells. *Mol Cell Endocrinol* **149**, 29-40, doi:10.1016/s0303-7207(99)00003-9 (1999).
- 28 Sowter, H. M. *et al.* Expression and localization of the vascular endothelial growth factor family in ovarian epithelial tumors. *Lab Invest* **77**, 607-614 (1997).
- 29 Niki, T. *et al.* Expression of vascular endothelial growth factors A, B, C, and D and their relationships to lymph node status in lung adenocarcinoma. *Clin Cancer Res* **6**, 2431-2439 (2000).
- 30 Mehrotra, D., Wu, J., Papangelis, I. & Chun, H. J. Endothelium as a gatekeeper of fatty acid transport. *Trends Endocrinol Metab* **25**, 99-106, doi:10.1016/j.tem.2013.11.001 (2014).
- 31 Jornayvaz, F. R. & Shulman, G. I. Regulation of mitochondrial biogenesis. *Essays Biochem* **47**, 69-84, doi:10.1042/bse0470069 (2010).
- 32 Mehlem, A. *et al.* PGC-1alpha Coordinates Mitochondrial Respiratory Capacity and Muscular Fatty Acid Uptake via Regulation of VEGF-B. *Diabetes* **65**, 861-873, doi:10.2337/db15-1231 (2016).
- 33 Enholm, B. *et al.* Comparison of VEGF, VEGF-B, VEGF-C and Ang-1 mRNA regulation by serum, growth factors, oncoproteins and hypoxia. *Oncogene* **14**, 2475-2483, doi:10.1038/sj.onc.1201090 (1997).
- 34 Singh, N. K., Hansen, D. E., 3rd, Kundumani-Sridharan, V. & Rao, G. N. Both Kdr and Flt1 play a vital role in hypoxia-induced Src-PLD1-PKCgamma-cPLA(2) activation and retinal neovascularization. *Blood* **121**, 1911-1923, doi:10.1182/blood-2012-03-419234 (2013).
- 35 Li, X. & Eriksson, U. Novel VEGF family members: VEGF-B, VEGF-C and VEGF-D. *The international journal of biochemistry & cell biology* **33**, 421-426 (2001).
- 36 Nash, A. D., Baca, M., Wright, C. & Scotney, P. D. The biology of vascular endothelial growth factor-B (VEGF-B). *Pulm Pharmacol Ther* **19**, 61-69, doi:10.1016/j.pupt.2005.02.007 (2006).
- 37 Aase, K. *et al.* Vascular endothelial growth factor-B-deficient mice display an atrial conduction defect. *Circulation* **104**, 358-364 (2001).
- 38 Bellomo, D. *et al.* Mice lacking the vascular endothelial growth factor-B gene (Vegfb) have smaller hearts, dysfunctional coronary vasculature, and impaired recovery from cardiac ischemia. *Circ Res* **86**, E29-35 (2000).
- 39 Reichelt, M. *et al.* Vascular endothelial growth factor-B and retinal vascular development in the mouse. *Clin Exp Ophthalmol* **31**, 61-65 (2003).
- 40 Louzier, V. *et al.* Role of VEGF-B in the lung during development of chronic hypoxic pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol* **284**, L926-937, doi:10.1152/ajplung.00247.2002 (2003).
- 41 Li, X. *et al.* Reevaluation of the role of VEGF-B suggests a restricted role in the revascularization of the ischemic myocardium. *Arteriosclerosis, thrombosis, and vascular biology* **28**, 1614-1620, doi:10.1161/ATVBAHA.107.158725 (2008).



- 42 Karpanen, T. *et al.* Overexpression of vascular endothelial growth factor-B in mouse heart alters cardiac lipid metabolism and induces myocardial hypertrophy. *Circ Res* **103**, 1018-1026, doi:10.1161/CIRCRESAHA.108.178459 (2008).
- 43 Mould, A. W. *et al.* Transgenic overexpression of vascular endothelial growth factor-B isoforms by endothelial cells potentiates postnatal vessel growth in vivo and in vitro. *Circ Res* **97**, e60-70, doi:10.1161/01.RES.0000182631.33638.77 (2005).
- 44 Detmar, M. *et al.* Increased microvascular density and enhanced leukocyte rolling and adhesion in the skin of VEGF transgenic mice. *J Invest Dermatol* **111**, 1-6, doi:10.1046/j.1523-1747.1998.00262.x (1998).
- 45 Larcher, F., Murillas, R., Bolontrade, M., Conti, C. J. & Jorcano, J. L. VEGF/VPF overexpression in skin of transgenic mice induces angiogenesis, vascular hyperpermeability and accelerated tumor development. *Oncogene* **17**, 303-311, doi:10.1038/sj.onc.1201928 (1998).
- 46 Xia, Y. P. *et al.* Transgenic delivery of VEGF to mouse skin leads to an inflammatory condition resembling human psoriasis. *Blood* **102**, 161-168, doi:10.1182/blood-2002-12-3793 (2003).
- 47 Jeltsch, M. *et al.* Hyperplasia of lymphatic vessels in VEGF-C transgenic mice. *Science* **276**, 1423-1425 (1997).
- 48 Karkkainen, A. M. *et al.* Vascular endothelial growth factor-D transgenic mice show enhanced blood capillary density, improved postischemic muscle regeneration, and increased susceptibility to tumor formation. *Blood* **113**, 4468-4475, doi:10.1182/blood-2008-07-171108 (2009).
- 49 Kiba, A., Sagara, H., Hara, T. & Shibuya, M. VEGFR-2-specific ligand VEGF-E induces non-edematous hyper-vascularization in mice. *Biochemical and biophysical research communications* **301**, 371-377 (2003).
- 50 Bhardwaj, S. *et al.* Angiogenic responses of vascular endothelial growth factors in periadventitial tissue. *Hum Gene Ther* **14**, 1451-1462, doi:10.1089/104303403769211664 (2003).
- 51 Rissanen, T. T. *et al.* VEGF-D is the strongest angiogenic and lymphangiogenic effector among VEGFs delivered into skeletal muscle via adenoviruses. *Circ Res* **92**, 1098-1106, doi:10.1161/01.RES.0000073584.46059.E3 (2003).
- 52 Zhong, X. *et al.* Vascular endothelial growth factor-B gene transfer exacerbates retinal and choroidal neovascularization and vasopermeability without promoting inflammation. *Mol Vis* **17**, 492-507 (2011).
- 53 Robciuc, M. R. *et al.* VEGFB/VEGFR1-Induced Expansion of Adipose Vasculature Counteracts Obesity and Related Metabolic Complications. *Cell Metab* **23**, 712-724, doi:10.1016/j.cmet.2016.03.004 (2016).
- 54 Lahteenvuo, J. E. *et al.* Vascular endothelial growth factor-B induces myocardium-specific angiogenesis and arteriogenesis via vascular endothelial growth factor receptor-1- and neuropilin receptor-1-dependent mechanisms. *Circulation* **119**, 845-856, doi:10.1161/CIRCULATIONAHA.108.816454 (2009).
- 55 Hagberg, C. E. *et al.* Targeting VEGF-B as a novel treatment for insulin resistance and type 2 diabetes. *Nature* **490**, 426-430, doi:10.1038/nature11464 (2012).
- 56 Kivela, R. *et al.* VEGF-B-induced vascular growth leads to metabolic reprogramming and ischemia resistance in the heart. *EMBO Mol Med* **6**, 307-321, doi:10.1002/emmm.201303147 (2014).
- 57 Falkevall, A. *et al.* Reducing VEGF-B Signaling Ameliorates Renal Lipotoxicity and Protects against Diabetic Kidney Disease. *Cell Metab* **25**, 713-726, doi:10.1016/j.cmet.2017.01.004 (2017).
- 58 Mathis, D. Immunological goings-on in visceral adipose tissue. *Cell metabolism* **17**, 851-859, doi:10.1016/j.cmet.2013.05.008 (2013).
- 59 Lu, X. *et al.* Resistance to obesity by repression of VEGF gene expression through induction of brown-like adipocyte differentiation. *Endocrinology* **153**, 3123-3132, doi:10.1210/en.2012-1151 (2012).
- 60 Bartelt, A. *et al.* Brown adipose tissue activity controls triglyceride clearance. *Nature medicine* **17**, 200-205, doi:10.1038/nm.2297 (2011).
- 61 Langerhans, P. Ueber die Nerven der menschlichen Haut. *Archiv für pathologische Anatomie und Physiologie und für klinische Medizin* **44**, 325-337, doi:10.1007/BF01959006 (1868).

- 62 Eberhard, D., Kragl, M. & Lammert, E. 'Giving and taking': endothelial and beta-cells in the islets of Langerhans. *Trends in endocrinology and metabolism: TEM* **21**, 457-463, doi:10.1016/j.tem.2010.03.003 (2010).
- 63 Kim, A. *et al.* Islet architecture: A comparative study. *Islets* **1**, 129-136, doi:10.4161/isl.1.2.9480 (2009).
- 64 Ionescu-Tirgoviste, C. *et al.* A 3D map of the islet routes throughout the healthy human pancreas. *Scientific reports* **5**, 14634, doi:10.1038/srep14634 (2015).
- 65 Henquin, J. C. The dual control of insulin secretion by glucose involves triggering and amplifying pathways in beta-cells. *Diabetes Res Clin Pract* **93 Suppl 1**, S27-31, doi:10.1016/S0168-8227(11)70010-9 (2011).
- 66 Curry, D. L., Bennett, L. L. & Grodsky, G. M. Dynamics of insulin secretion by the perfused rat pancreas. *Endocrinology* **83**, 572-584, doi:10.1210/endo-83-3-572 (1968).
- 67 Reinert, R. B. *et al.* Vascular endothelial growth factor- $\alpha$  and islet vascularization are necessary in developing, but not adult, pancreatic islets. *Diabetes* **62**, 4154-4164, doi:10.2337/db13-0071 (2013).
- 68 Brissova, M. *et al.* Pancreatic islet production of vascular endothelial growth factor- $\alpha$  is essential for islet vascularization, revascularization, and function. *Diabetes* **55**, 2974-2985, doi:10.2337/db06-0690 (2006).
- 69 Konstantinova, I. & Lammert, E. Microvascular development: learning from pancreatic islets. *BioEssays : news and reviews in molecular, cellular and developmental biology* **26**, 1069-1075, doi:10.1002/bies.20105 (2004).
- 70 Lammert, E. *et al.* Role of VEGF-A in vascularization of pancreatic islets. *Current biology : CB* **13**, 1070-1074 (2003).
- 71 Cabrera, O. *et al.* The unique cytoarchitecture of human pancreatic islets has implications for islet cell function. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 2334-2339, doi:10.1073/pnas.0510790103 (2006).
- 72 Olsson, R. & Carlsson, P. O. The pancreatic islet endothelial cell: emerging roles in islet function and disease. *The international journal of biochemistry & cell biology* **38**, 710-714 (2006).
- 73 Nyman, L. R. *et al.* Real-time, multidimensional in vivo imaging used to investigate blood flow in mouse pancreatic islets. *The Journal of clinical investigation* **118**, 3790-3797, doi:10.1172/JCI36209 (2008).
- 74 Ballian, N. & Brunicardi, F. C. Islet vasculature as a regulator of endocrine pancreas function. *World journal of surgery* **31**, 705-714, doi:10.1007/s00268-006-0719-8 (2007).
- 75 Jansson, L. The regulation of pancreatic islet blood flow. *Diabetes/metabolism reviews* **10**, 407-416 (1994).
- 76 Weir, G. C. & Bonner-Weir, S. Five stages of evolving beta-cell dysfunction during progression to diabetes. *Diabetes* **53 Suppl 3**, S16-21 (2004).
- 77 Carlsson, P. O., Jansson, L., Ostenson, C. G. & Kallskog, O. Islet capillary blood pressure increase mediated by hyperglycemia in NIDDM GK rats. *Diabetes* **46**, 947-952 (1997).
- 78 Schmidt, H. H., Warner, T. D., Ishii, K., Sheng, H. & Murad, F. Insulin secretion from pancreatic B cells caused by L-arginine-derived nitrogen oxides. *Science* **255**, 721-723 (1992).
- 79 Henningsson, R., Salehi, A. & Lundquist, I. Role of nitric oxide synthase isoforms in glucose-stimulated insulin release. *American journal of physiology. Cell physiology* **283**, C296-304, doi:10.1152/ajpcell.00537.2001 (2002).
- 80 Lacraz, G. *et al.* Islet endothelial activation and oxidative stress gene expression is reduced by IL-1Ra treatment in the type 2 diabetic GK rat. *PloS one* **4**, e6963, doi:10.1371/journal.pone.0006963 (2009).
- 81 Li, X. *et al.* Islet microvasculature in islet hyperplasia and failure in a model of type 2 diabetes. *Diabetes* **55**, 2965-2973, doi:10.2337/db06-0733 (2006).
- 82 Staiger, K. *et al.* Saturated, but not unsaturated, fatty acids induce apoptosis of human coronary artery endothelial cells via nuclear factor- $\kappa$ B activation. *Diabetes* **55**, 3121-3126, doi:10.2337/db06-0188 (2006).
- 83 Glatz, J. F., Luiken, J. J. & Bonen, A. Membrane fatty acid transporters as regulators of lipid metabolism: implications for metabolic disease. *Physiological reviews* **90**, 367-417, doi:10.1152/physrev.00003.2009 (2010).

- 84 Eiselein, L., Wilson, D. W., Lame, M. W. & Rutledge, J. C. Lipolysis products from triglyceride-rich lipoproteins increase endothelial permeability, perturb zonula occludens-1 and F-actin, and induce apoptosis. *American journal of physiology. Heart and circulatory physiology* **292**, H2745-2753, doi:10.1152/ajpheart.00686.2006 (2007).
- 85 Kazantzis, M. & Stahl, A. Fatty acid transport proteins, implications in physiology and disease. *Biochimica et biophysica acta* **1821**, 852-857, doi:10.1016/j.bbalip.2011.09.010 (2012).
- 86 Bharadwaj, K. G. *et al.* Chylomicron- and VLDL-derived lipids enter the heart through different pathways: in vivo evidence for receptor- and non-receptor-mediated fatty acid uptake. *The Journal of biological chemistry* **285**, 37976-37986, doi:10.1074/jbc.M110.174458 (2010).
- 87 Silverstein, R. L. & Febbraio, M. CD36, a scavenger receptor involved in immunity, metabolism, angiogenesis, and behavior. *Science signaling* **2**, re3, doi:10.1126/scisignal.272re3 (2009).
- 88 Asch, A. S., Barnwell, J., Silverstein, R. L. & Nachman, R. L. Isolation of the thrombospondin membrane receptor. *The Journal of clinical investigation* **79**, 1054-1061, doi:10.1172/JCI112918 (1987).
- 89 Hoebe, K. *et al.* CD36 is a sensor of diacylglycerides. *Nature* **433**, 523-527, doi:10.1038/nature03253 (2005).
- 90 Martin, C. *et al.* CD36 as a lipid sensor. *Physiology & behavior* **105**, 36-42, doi:10.1016/j.physbeh.2011.02.029 (2011).
- 91 Podrez, E. A. *et al.* Macrophage scavenger receptor CD36 is the major receptor for LDL modified by monocyte-generated reactive nitrogen species. *The Journal of clinical investigation* **105**, 1095-1108, doi:10.1172/JCI8574 (2000).
- 92 Bastie, C. C., Hajri, T., Drover, V. A., Grimaldi, P. A. & Abumrad, N. A. CD36 in myocytes channels fatty acids to a lipase-accessible triglyceride pool that is related to cell lipid and insulin responsiveness. *Diabetes* **53**, 2209-2216 (2004).
- 93 Carley, A. N. & Kleinfeld, A. M. Fatty acid (FFA) transport in cardiomyocytes revealed by imaging unbound FFA is mediated by an FFA pump modulated by the CD36 protein. *The Journal of biological chemistry* **286**, 4589-4597, doi:10.1074/jbc.M110.182162 (2011).
- 94 Rosen, E. D. & Spiegelman, B. M. Peroxisome proliferator-activated receptor gamma ligands and atherosclerosis: ending the heartache. *The Journal of clinical investigation* **106**, 629-631, doi:10.1172/JCI10909 (2000).
- 95 Tontonoz, P., Nagy, L., Alvarez, J. G., Thomazy, V. A. & Evans, R. M. PPARgamma promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* **93**, 241-252 (1998).
- 96 Kanda, T. *et al.* PPARgamma in the endothelium regulates metabolic responses to high-fat diet in mice. *The Journal of clinical investigation* **119**, 110-124, doi:10.1172/JCI36233 (2009).
- 97 Kuda, O. *et al.* Sulfo-N-succinimidyl oleate (SSO) inhibits fatty acid uptake and signaling for intracellular calcium via binding CD36 lysine 164: SSO also inhibits oxidized low density lipoprotein uptake by macrophages. *The Journal of biological chemistry* **288**, 15547-15555, doi:10.1074/jbc.M113.473298 (2013).
- 98 Doege, H. & Stahl, A. Protein-mediated fatty acid uptake: novel insights from in vivo models. *Physiology* **21**, 259-268, doi:10.1152/physiol.00014.2006 (2006).
- 99 Hirsch, D., Stahl, A. & Lodish, H. F. A family of fatty acid transporters conserved from mycobacterium to man. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 8625-8629 (1998).
- 100 Schaffer, J. E. & Lodish, H. F. Expression cloning and characterization of a novel adipocyte long chain fatty acid transport protein. *Cell* **79**, 427-436 (1994).
- 101 Anderson, C. M. & Stahl, A. SLC27 fatty acid transport proteins. *Mol Aspects Med* **34**, 516-528, doi:10.1016/j.mam.2012.07.010 (2013).
- 102 Steinberg, S. J., Wang, S. J., McGuinness, M. C. & Watkins, P. A. Human liver-specific very-long-chain acyl-coenzyme A synthetase: cDNA cloning and characterization of a second enzymatically active protein. *Molecular genetics and metabolism* **68**, 32-42, doi:10.1006/mgme.1999.2883 (1999).
- 103 Pei, Z. *et al.* Mouse very long-chain Acyl-CoA synthetase 3/fatty acid transport protein 3 catalyzes fatty acid activation but not fatty acid transport in MA-10 cells. *The Journal of biological chemistry* **279**, 54454-54462, doi:10.1074/jbc.M410091200 (2004).

- 104 Stahl, A. *et al.* Identification of the major intestinal fatty acid transport protein. *Molecular cell* **4**, 299-308 (1999).
- 105 Sandoval, A. *et al.* Fatty acid transport and activation and the expression patterns of genes involved in fatty acid trafficking. *Archives of biochemistry and biophysics* **477**, 363-371, doi:10.1016/j.abb.2008.06.010 (2008).
- 106 DiRusso, C. C. *et al.* Comparative biochemical studies of the murine fatty acid transport proteins (FATP) expressed in yeast. *The Journal of biological chemistry* **280**, 16829-16837, doi:10.1074/jbc.M409598200 (2005).
- 107 Milger, K. *et al.* Cellular uptake of fatty acids driven by the ER-localized acyl-CoA synthetase FATP4. *Journal of cell science* **119**, 4678-4688, doi:10.1242/jcs.03280 (2006).
- 108 Stahl, A., Gimeno, R. E., Tartaglia, L. A. & Lodish, H. F. Fatty acid transport proteins: a current view of a growing family. *Trends in endocrinology and metabolism: TEM* **12**, 266-273 (2001).
- 109 Shimabukuro, M., Zhou, Y. T., Levi, M. & Unger, R. H. Fatty acid-induced beta cell apoptosis: a link between obesity and diabetes. *Proc Natl Acad Sci U S A* **95**, 2498-2502, doi:10.1073/pnas.95.5.2498 (1998).
- 110 Moulle, V. S., Ghislain, J. & Poitout, V. Nutrient regulation of pancreatic beta-cell proliferation. *Biochimie* **143**, 10-17, doi:10.1016/j.biochi.2017.09.017 (2017).
- 111 Acosta-Montano, P. & Garcia-Gonzalez, V. Effects of Dietary Fatty Acids in Pancreatic Beta Cell Metabolism, Implications in Homeostasis. *Nutrients* **10**, doi:10.3390/nu10040393 (2018).
- 112 Prentki, M. & Madiraju, S. R. Glycerolipid/free fatty acid cycle and islet beta-cell function in health, obesity and diabetes. *Mol Cell Endocrinol* **353**, 88-100, doi:10.1016/j.mce.2011.11.004 (2012).
- 113 Cruz, W. S., Kwon, G., Marshall, C. A., McDaniel, M. L. & Semenkovich, C. F. Glucose and insulin stimulate heparin-releasable lipoprotein lipase activity in mouse islets and INS-1 cells. A potential link between insulin resistance and beta-cell dysfunction. *J Biol Chem* **276**, 12162-12168, doi:10.1074/jbc.M010707200 (2001).
- 114 Pappan, K. L. *et al.* Pancreatic beta-cell lipoprotein lipase independently regulates islet glucose metabolism and normal insulin secretion. *J Biol Chem* **280**, 9023-9029, doi:10.1074/jbc.M409706200 (2005).
- 115 Berk, P. D. *et al.* Plasma membrane fatty acid-binding protein and mitochondrial glutamic-oxaloacetic transaminase of rat liver are related. *Proc Natl Acad Sci U S A* **87**, 3484-3488, doi:10.1073/pnas.87.9.3484 (1990).
- 116 Stremmel, W., Strohmeyer, G., Borchard, F., Kochwa, S. & Berk, P. D. Isolation and partial characterization of a fatty acid binding protein in rat liver plasma membranes. *Proc Natl Acad Sci U S A* **82**, 4-8, doi:10.1073/pnas.82.1.4 (1985).
- 117 Furuhashi, M. & Hotamisligil, G. S. Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets. *Nat Rev Drug Discov* **7**, 489-503, doi:10.1038/nrd2589 (2008).
- 118 Noshmehr, H. *et al.* Fatty acid translocase (FAT/CD36) is localized on insulin-containing granules in human pancreatic beta-cells and mediates fatty acid effects on insulin secretion. *Diabetes* **54**, 472-481, doi:10.2337/diabetes.54.2.472 (2005).
- 119 Wallin, T. *et al.* Facilitation of fatty acid uptake by CD36 in insulin-producing cells reduces fatty-acid-induced insulin secretion and glucose regulation of fatty acid oxidation. *Biochim Biophys Acta* **1801**, 191-197, doi:10.1016/j.bbalip.2009.11.002 (2010).
- 120 Vangaveti, V., Shashidhar, V., Jarrod, G., Baune, B. T. & Kennedy, R. L. Free fatty acid receptors: emerging targets for treatment of diabetes and its complications. *Ther Adv Endocrinol Metab* **1**, 165-175, doi:10.1177/2042018810381066 (2010).
- 121 Itoh, Y. *et al.* Free fatty acids regulate insulin secretion from pancreatic beta cells through GPR40. *Nature* **422**, 173-176, doi:10.1038/nature01478 (2003).
- 122 Miyauchi, S., Hirasawa, A., Ichimura, A., Hara, T. & Tsujimoto, G. New frontiers in gut nutrient sensor research: free fatty acid sensing in the gastrointestinal tract. *J Pharmacol Sci* **112**, 19-24, doi:10.1254/jphs.09r09fm (2010).
- 123 Tuo, Y. *et al.* Long-term in vitro treatment of INS-1 rat pancreatic beta-cells by unsaturated free fatty acids protects cells against gluco- and lipotoxicities via activation of GPR40 receptors. *Clin Exp Pharmacol Physiol* **39**, 423-428, doi:10.1111/j.1440-1681.2012.05691.x (2012).

- 124 Nagasumi, K. *et al.* Overexpression of GPR40 in pancreatic beta-cells augments glucose-stimulated insulin secretion and improves glucose tolerance in normal and diabetic mice. *Diabetes* **58**, 1067-1076, doi:10.2337/db08-1233 (2009).
- 125 Wagner, R. *et al.* Reevaluation of fatty acid receptor 1 as a drug target for the stimulation of insulin secretion in humans. *Diabetes* **62**, 2106-2111, doi:10.2337/db12-1249 (2013).
- 126 Steneberg, P., Rubins, N., Bartoov-Shifman, R., Walker, M. D. & Edlund, H. The FFA receptor GPR40 links hyperinsulinemia, hepatic steatosis, and impaired glucose homeostasis in mouse. *Cell Metab* **1**, 245-258, doi:10.1016/j.cmet.2005.03.007 (2005).
- 127 Kristinsson, H., Smith, D. M., Bergsten, P. & Sargsyan, E. FFAR1 is involved in both the acute and chronic effects of palmitate on insulin secretion. *Endocrinology* **154**, 4078-4088, doi:10.1210/en.2013-1352 (2013).
- 128 Hirasawa, A. *et al.* Free fatty acids regulate gut incretin glucagon-like peptide-1 secretion through GPR120. *Nat Med* **11**, 90-94, doi:10.1038/nm1168 (2005).
- 129 Katsuma, S. *et al.* Free fatty acids inhibit serum deprivation-induced apoptosis through GPR120 in a murine enteroendocrine cell line STC-1. *J Biol Chem* **280**, 19507-19515, doi:10.1074/jbc.M412385200 (2005).
- 130 Ichimura, A. *et al.* Dysfunction of lipid sensor GPR120 leads to obesity in both mouse and human. *Nature* **483**, 350-354, doi:10.1038/nature10798 (2012).
- 131 Tang, C. *et al.* Loss of FFA2 and FFA3 increases insulin secretion and improves glucose tolerance in type 2 diabetes. *Nat Med* **21**, 173-177, doi:10.1038/nm.3779 (2015).
- 132 Tolhurst, G. *et al.* Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein-coupled receptor FFAR2. *Diabetes* **61**, 364-371, doi:10.2337/db11-1019 (2012).
- 133 Grygiel-Gorniak, B. Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications--a review. *Nutr J* **13**, 17, doi:10.1186/1475-2891-13-17 (2014).
- 134 Popescu, I. R. *et al.* The nuclear receptor FXR is expressed in pancreatic beta-cells and protects human islets from lipotoxicity. *FEBS Lett* **584**, 2845-2851, doi:10.1016/j.febslet.2010.04.068 (2010).
- 135 Mokdad, A. H. *et al.* Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *Jama* **289**, 76-79 (2003).
- 136 Boucher, A. *et al.* Biochemical mechanism of lipid-induced impairment of glucose-stimulated insulin secretion and reversal with a malate analogue. *The Journal of biological chemistry* **279**, 27263-27271, doi:10.1074/jbc.M401167200 (2004).
- 137 Arner, P. & Ryden, M. Fatty Acids, Obesity and Insulin Resistance. *Obesity facts* **8**, 147-155, doi:10.1159/000381224 (2015).
- 138 Fonseca, S. G., Gromada, J. & Urano, F. Endoplasmic reticulum stress and pancreatic beta-cell death. *Trends in endocrinology and metabolism: TEM* **22**, 266-274, doi:10.1016/j.tem.2011.02.008 (2011).
- 139 Biden, T. J., Boslem, E., Chu, K. Y. & Sue, N. Lipotoxic endoplasmic reticulum stress, beta cell failure, and type 2 diabetes mellitus. *Trends in endocrinology and metabolism: TEM* **25**, 389-398, doi:10.1016/j.tem.2014.02.003 (2014).
- 140 Hara, T. *et al.* Calcium efflux from the endoplasmic reticulum leads to beta-cell death. *Endocrinology* **155**, 758-768, doi:10.1210/en.2013-1519 (2014).
- 141 Boslem, E. *et al.* Alteration of endoplasmic reticulum lipid rafts contributes to lipotoxicity in pancreatic beta-cells. *The Journal of biological chemistry* **288**, 26569-26582, doi:10.1074/jbc.M113.489310 (2013).
- 142 Back, S. H., Kang, S. W., Han, J. & Chung, H. T. Endoplasmic reticulum stress in the beta-cell pathogenesis of type 2 diabetes. *Experimental diabetes research* **2012**, 618396, doi:10.1155/2012/618396 (2012).
- 143 Cunha, D. A. *et al.* Initiation and execution of lipotoxic ER stress in pancreatic beta-cells. *Journal of cell science* **121**, 2308-2318, doi:10.1242/jcs.026062 (2008).
- 144 Graciano, M. F., Valle, M. M., Kowluru, A., Curi, R. & Carpinelli, A. R. Regulation of insulin secretion and reactive oxygen species production by free fatty acids in pancreatic islets. *Islets* **3**, 213-223 (2011).
- 145 Barlow, J. & Affourtit, C. Novel insights into pancreatic beta-cell glucolipotoxicity from real-time functional analysis of mitochondrial energy metabolism in INS-1E insulinoma cells. *The Biochemical journal* **456**, 417-426, doi:10.1042/BJ20131002 (2013).

- 146 Karunakaran, U. & Park, K. G. A systematic review of oxidative stress and safety of antioxidants in diabetes: focus on islets and their defense. *Diabetes & metabolism journal* **37**, 106-112, doi:10.4093/dmj.2013.37.2.106 (2013).
- 147 Maechler, P. & Wollheim, C. B. Mitochondrial function in normal and diabetic beta-cells. *Nature* **414**, 807-812, doi:10.1038/414807a (2001).
- 148 Sharma, R. B. & Alonso, L. C. Lipotoxicity in the pancreatic beta cell: not just survival and function, but proliferation as well? *Current diabetes reports* **14**, 492, doi:10.1007/s11892-014-0492-2 (2014).
- 149 Watada, H. & Fujitani, Y. Minireview: Autophagy in pancreatic beta-cells and its implication in diabetes. *Molecular endocrinology* **29**, 338-348, doi:10.1210/me.2014-1367 (2015).
- 150 Ebato, C. *et al.* Autophagy is important in islet homeostasis and compensatory increase of beta cell mass in response to high-fat diet. *Cell metabolism* **8**, 325-332, doi:10.1016/j.cmet.2008.08.009 (2008).
- 151 Masini, M. *et al.* Autophagy in human type 2 diabetes pancreatic beta cells. *Diabetologia* **52**, 1083-1086, doi:10.1007/s00125-009-1347-2 (2009).
- 152 Las, G., Serada, S. B., Wikstrom, J. D., Twig, G. & Shirihai, O. S. Fatty acids suppress autophagic turnover in beta-cells. *The Journal of biological chemistry* **286**, 42534-42544, doi:10.1074/jbc.M111.242412 (2011).
- 153 Kahn, S. E., Zraika, S., Utzschneider, K. M. & Hull, R. L. The beta cell lesion in type 2 diabetes: there has to be a primary functional abnormality. *Diabetologia* **52**, 1003-1012, doi:10.1007/s00125-009-1321-z (2009).
- 154 Rhodes, C. J. Type 2 diabetes-a matter of beta-cell life and death? *Science* **307**, 380-384, doi:10.1126/science.1104345 (2005).
- 155 Halban, P. A. *et al.* beta-cell failure in type 2 diabetes: postulated mechanisms and prospects for prevention and treatment. *Diabetes Care* **37**, 1751-1758, doi:10.2337/dc14-0396 (2014).
- 156 Barker, C. J., Leibiger, I. B. & Berggren, P. O. The pancreatic islet as a signaling hub. *Adv Biol Regul* **53**, 156-163, doi:10.1016/j.jbior.2012.09.011 (2013).
- 157 Leibiger, I. B. & Berggren, P. O. Intraocular in vivo imaging of pancreatic islet cell physiology/pathology. *Mol Metab* **6**, 1002-1009, doi:10.1016/j.molmet.2017.03.014 (2017).
- 158 Saudek, F., Brogren, C. H. & Manohar, S. Imaging the Beta-cell mass: why and how. *Rev Diabet Stud* **5**, 6-12, doi:10.1900/RDS.2008.5.6 (2008).
- 159 Yang, S. N. & Berggren, P. O. The eye as a novel imaging site in diabetes research. *Pharmacol Ther* **197**, 103-121, doi:10.1016/j.pharmthera.2019.01.005 (2019).
- 160 Speier, S. *et al.* Noninvasive in vivo imaging of pancreatic islet cell biology. *Nat Med* **14**, 574-578, doi:10.1038/nm1701 (2008).
- 161 Speier, S. *et al.* Noninvasive high-resolution in vivo imaging of cell biology in the anterior chamber of the mouse eye. *Nat Protoc* **3**, 1278-1286, doi:10.1038/nprot.2008.118 (2008).
- 162 Abdulreda, M. H., Caicedo, A. & Berggren, P. O. Transplantation into the anterior chamber of the eye for longitudinal, non-invasive in vivo imaging with single-cell resolution in real-time. *J Vis Exp*, e50466, doi:10.3791/50466 (2013).
- 163 Almaca, J. *et al.* Young capillary vessels rejuvenate aged pancreatic islets. *Proc Natl Acad Sci U S A* **111**, 17612-17617, doi:10.1073/pnas.1414053111 (2014).
- 164 Nyqvist, D. *et al.* Donor islet endothelial cells in pancreatic islet revascularization. *Diabetes* **60**, 2571-2577, doi:10.2337/db10-1711 (2011).
- 165 Rodriguez-Diaz, R. *et al.* Noninvasive in vivo model demonstrating the effects of autonomic innervation on pancreatic islet function. *Proc Natl Acad Sci U S A* **109**, 21456-21461, doi:10.1073/pnas.1211659110 (2012).
- 166 Ali, Y. *et al.* The anterior chamber of the eye is a transplantation site that supports and enables visualisation of beta cell development in mice. *Diabetologia* **59**, 1007-1011, doi:10.1007/s00125-016-3883-x (2016).
- 167 Chen, C. *et al.* Alterations in beta-Cell Calcium Dynamics and Efficacy Outweigh Islet Mass Adaptation in Compensation of Insulin Resistance and Prediabetes Onset. *Diabetes* **65**, 2676-2685, doi:10.2337/db15-1718 (2016).
- 168 Paschen, M. *et al.* Non-invasive cell type selective in vivo monitoring of insulin resistance dynamics. *Sci Rep* **6**, 21448, doi:10.1038/srep21448 (2016).

169 Paschen, M. *et al.* Diet-induced beta-cell insulin resistance results in reversible loss of functional  
beta-cell mass. *FASEB J* **33**, 204-218, doi:10.1096/fj.201800826R (2019).

170 Hanahan, D. Heritable formation of pancreatic beta-cell tumours in transgenic mice expressing  
recombinant insulin/simian virus 40 oncogenes. *Nature* **315**, 115-122, doi:10.1038/315115a0  
(1985).

171 Bill, R. & Christofori, G. The Rip1Tag2 Transgenic Mouse Model. *Methods Mol Biol* **1464**, 151-  
161, doi:10.1007/978-1-4939-3999-2\_14 (2016).

172 Nusse, R. *et al.* A new nomenclature for int-1 and related genes: the Wnt gene family. *Cell* **64**,  
231, doi:10.1016/0092-8674(91)90633-a (1991).

173 Nusse, R. & Varmus, H. E. Wnt genes. *Cell* **69**, 1073-1087, doi:10.1016/0092-8674(92)90630-u  
(1992).

174 Nusse, R. Wnt signaling in disease and in development. *Cell Res* **15**, 28-32,  
doi:10.1038/sj.cr.7290260 (2005).

175 Katoh, M. WNT/PCP signaling pathway and human cancer (review). *Oncol Rep* **14**, 1583-1588  
(2005).

176 Sugimura, R. & Li, L. Noncanonical Wnt signaling in vertebrate development, stem cells, and  
diseases. *Birth Defects Res C Embryo Today* **90**, 243-256, doi:10.1002/bdrc.20195 (2010).

177 Kohn, A. D. & Moon, R. T. Wnt and calcium signaling: beta-catenin-independent pathways.  
*Cell Calcium* **38**, 439-446, doi:10.1016/j.ceca.2005.06.022 (2005).

178 De, A. Wnt/Ca<sup>2+</sup> signaling pathway: a brief overview. *Acta Biochim Biophys Sin (Shanghai)* **43**,  
745-756, doi:10.1093/abbs/gmr079 (2011).

179 Clevers, H. Wnt/beta-catenin signaling in development and disease. *Cell* **127**, 469-480,  
doi:10.1016/j.cell.2006.10.018 (2006).

180 Kimelman, D. & Xu, W. beta-catenin destruction complex: insights and questions from a  
structural perspective. *Oncogene* **25**, 7482-7491, doi:10.1038/sj.onc.1210055 (2006).

181 Aberle, H., Bauer, A., Stappert, J., Kispert, A. & Kemler, R. beta-catenin is a target for the  
ubiquitin-proteasome pathway. *EMBO J* **16**, 3797-3804, doi:10.1093/emboj/16.13.3797 (1997).

182 MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components, mechanisms,  
and diseases. *Dev Cell* **17**, 9-26, doi:10.1016/j.devcel.2009.06.016 (2009).

183 MacDonald, B. T. & He, X. Frizzled and LRP5/6 receptors for Wnt/beta-catenin signaling.  
*Cold Spring Harb Perspect Biol* **4**, doi:10.1101/cshperspect.a007880 (2012).

184 Soares, M. B. *et al.* RNA-mediated gene duplication: the rat preproinsulin I gene is a functional  
retropon. *Mol Cell Biol* **5**, 2090-2103, doi:10.1128/mcb.5.8.2090 (1985).

185 Wentworth, B. M., Schaefer, I. M., Villa-Komaroff, L. & Chirgwin, J. M. Characterization of  
the two nonallelic genes encoding mouse preproinsulin. *J Mol Evol* **23**, 305-312 (1986).

186 Andrali, S. S., Sampley, M. L., Vanderford, N. L. & Ozcan, S. Glucose regulation of insulin  
gene expression in pancreatic beta-cells. *Biochem J* **415**, 1-10, doi:10.1042/BJ20081029 (2008).

187 Matsuoka, T. A. *et al.* MafA regulates expression of genes important to islet beta-cell function.  
*Mol Endocrinol* **21**, 2764-2774, doi:10.1210/me.2007-0028 (2007).

188 Song, Y. D. *et al.* Islet cell differentiation in liver by combinatorial expression of transcription  
factors neurogenin-3, BETA2, and RIPE3b1. *Biochem Biophys Res Commun* **354**, 334-339,  
doi:10.1016/j.bbrc.2006.12.216 (2007).

189 Ohneda, K., Mirmira, R. G., Wang, J., Johnson, J. D. & German, M. S. The homeodomain of  
PDX-1 mediates multiple protein-protein interactions in the formation of a transcriptional  
activation complex on the insulin promoter. *Mol Cell Biol* **20**, 900-911,  
doi:10.1128/mcb.20.3.900-911.2000 (2000).

190 Valtat, B. *et al.* Fetal PGC-1alpha overexpression programs adult pancreatic beta-cell  
dysfunction. *Diabetes* **62**, 1206-1216, doi:10.2337/db12-0314 (2013).

191 Li, X. *et al.* VEGF-B: a survival, or an angiogenic factor? *Cell Adh Migr* **3**, 322-327,  
doi:10.4161/cam.3.4.9459 (2009).

192 Tran, K. A. *et al.* Endothelial beta-Catenin Signaling Is Required for Maintaining Adult Blood-  
Brain Barrier Integrity and Central Nervous System Homeostasis. *Circulation* **133**, 177-186,  
doi:10.1161/CIRCULATIONAHA.115.015982 (2016).

193 Phng, L. K. *et al.* Nrarp coordinates endothelial Notch and Wnt signaling to control vessel  
density in angiogenesis. *Dev Cell* **16**, 70-82, doi:10.1016/j.devcel.2008.12.009 (2009).

- 194 Corada, M. *et al.* The Wnt/beta-catenin pathway modulates vascular remodeling and specification by upregulating Dll4/Notch signaling. *Dev Cell* **18**, 938-949, doi:10.1016/j.devcel.2010.05.006 (2010).